

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2002 (03.01.2002)

PCT

(10) International Publication Number
WO 02/00856 A2

- (51) International Patent Classification⁷: C12N 9/00 M.; Ketelbuiserstraat 99, B-8810 Lichtervelde (BE).
GEYSENS, Steven, C., J.; Steenbeekstraaje 8, B-9770
Kruishoutem (BE).
- (21) International Application Number: PCT/IB01/01514
- (22) International Filing Date: 29 June 2001 (29.06.2001) (74) Agent: PUST, Detlev, A.; Hoffmann . Eitle, Arabellas-
trasse 4, 81925 München (DE).
- (25) Filing Language: English (81) Designated States (*national*): AU, CA, JP, KR, MX, NO.
- (26) Publication Language: English (84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).
- (30) Priority Data: 60/215,676 30 June 2000 (30.06.2000) US
Published:
— without international search report and to be republished
upon receipt of that report
- (71) Applicant: FLANDERS INTERUNIVERSITY INSTI-
TUTE FOR BIOTECHNOLOGY (VIB) [BE/BE]; Ri-
jvissestraat 120, B-9052 Zwijnaarde (BE).
For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.
- (72) Inventors: CONTRERAS, Roland; Molenstraat 53,
B-9820 Merelbeke (BE). CALLEWAERT, Nico, L.,



WO 02/00856 A2

(54) Title: PROTEIN GLYCOSYLATION MODIFICATION IN *PICHIA PASTORIS*

(57) Abstract: The present invention provides genetically engineered strains of *Pichia* capable of producing proteins with reduced glycosylation. In particular, the genetically engineered strains of the present invention are capable of expressing either or both of an α -1,2-mannosidase and glucosidase II. The genetically engineered strains of the present invention can be further modified such that the OCH1 gene is disrupted. Methods of producing glycoproteins with reduced glycosylation using such genetically engineered strains of *Pichia* are also provided.

Protein Glycosylation Modification in *Pichia Pastoris*

Field of the Invention

The present invention relates to methods and vectors useful for genetically
5 modifying the glycosylation process in methylotrophic yeast strains for the purpose of
producing glycoproteins with reduced glycosylation. The present invention further
relates to methylotrophic yeast strains generated using the present methods and
vectors, as well as glycoproteins produced from such genetically modified strains.

10 Background of the Invention

The methylotrophic yeasts including *Pichia pastoris* have been widely used
for production of recombinant proteins of commercial or medical importance.
However, production and medical applications of some therapeutic glycoproteins can
be hampered by the differences in the protein-linked carbohydrate biosynthesis
15 between these yeasts and the target organism such as a mammalian subject.

Protein N-glycosylation originates in the endoplasmic reticulum (ER),
where an N-linked oligosaccharide ($\text{Glc}_3\text{Man}_9\text{GlcNAc}_2$) assembled on dolichol (a lipid
carrier intermediate) is transferred to the appropriate Asn of a nascent protein. This is
an event common to all eukaryotic N-linked glycoproteins. The three glucose residues
20 and one specific α -1,2-linked mannose residue are removed by specific glucosidases
and an α -1,2-mannosidase in the ER, resulting in the core oligosaccharide structure,
 $\text{Man}_8\text{GlcNAc}_2$. The protein with this core sugar structure is transported to the Golgi
apparatus where the sugar moiety undergoes various modifications. There are
significant differences in the modifications of the sugar chain in the Golgi apparatus
25 between yeast and higher eukaryotes.

In mammalian cells, the modification of the sugar chain proceeds via 3
different pathways depending on the protein moiety to which it is added. That is, (1)
the core sugar chain does not change; (2) the core sugar chain is changed by adding
the N-acetylglucosamine-1-phosphate moiety (GlcNAc-1-P) in UDP-N-acetyl

glucosamine (UDP-GlcNAc) to the 6-position of mannose in the core sugar chain, followed by removing the GlcNAc moiety to form an acidic sugar chain in the glycoprotein; or (3) the core sugar chain is first converted into $\text{Man}_5\text{GlcNAc}_2$ by removing 3 mannose residues with mannosidase I; $\text{Man}_5\text{GlcNAc}_2$ is further modified
5 by adding GlcNAc and removing 2 more mannose residues, followed by sequentially adding GlcNAc, galactose (Gal), and N-acetylneuraminic acid (also called sialic acid (NeuNAc)) to form various hybrid or complex sugar chains (R. Kornfeld and S. Kornfeld, *Ann. Rev. Biochem.* 54: 631-664, 1985; Chiba et al *J. Biol. Chem.* 273: 26298-26304, 1998).

10 In yeast, the modification of the sugar chain in the Golgi involves a series of additions of mannose residues by different mannosyltransferases ("outer chain" glycosylation). The structure of the outer chain glycosylation is specific to the organisms, typically with more than 50 mannose residues in *S. cerevisiae*, and most commonly with structures smaller than $\text{Man}_{14}\text{GlcNAc}_2$ in *Pichia pastoris*. This yeast-
15 specific outer chain glycosylation of the high mannose type is also denoted hyperglycosylation.

Hyperglycosylation is often undesired since it leads to heterogeneity of a recombinant protein product in both carbohydrate composition and molecular weight, which may complicate the protein purification. The specific activity (units/weight) of
20 hyperglycosylated enzymes may be lowered by the increased portion of carbohydrate. In addition, the outer chain glycosylation is strongly immunogenic which is undesirable in a therapeutic application. Moreover, the large outer chain sugar can mask the immunogenic determinants of a therapeutic protein. For example, the influenza neuraminidase (NA) expressed in *P. pastoris* is glycosylated with N-glycans
25 containing up to 30-40 mannose residues. The hyperglycosylated NA has a reduced immunogenicity in mice, as the variable and immunodominant surface loops on top of the NA molecule are masked by the N-glycans (Martinet et al. *Eur J. Biochem.* 247: 332-338, 1997).

Therefore, it is desirable to genetically engineer methylotrophic yeast strains in which glycosylation of proteins can be manipulated and from which recombinant proteins can be produced that would not be compromised in structure or function by large N-glycan side chains.

5

Summary of the Invention

The present invention is directed to methods and vectors useful for genetically modifying the glycosylation process in methylotrophic yeast strains to produce glycoproteins with reduced glycosylation. Methylotrophic yeast strains
10 generated using the present methods and vectors, as well as glycoproteins produced from such genetically modified strains are also provided.

In one embodiment, the present invention provides vectors useful for making genetically engineered methylotrophic yeast strains which are capable of producing glycoproteins with reduced glycosylation.

15 In one aspect, the present invention provides "knock-in" vectors which are capable of expressing in a methylotrophic yeast strain one or more proteins whose enzymatic activities lead to a reduction of glycosylation in glycoproteins produced by the methylotrophic yeast strain.

In a preferred embodiment, the knock-in vectors of the present invention
20 include a nucleotide sequence coding for an α -1,2-mannosidase or a functional part thereof and are capable of expressing the α -1,2-mannosidase or the functional part in a methylotrophic yeast strain. A preferred nucleotide sequence is a nucleotide sequence encoding the α -1,2-mannosidase of a fungal species, and more preferably, *Trichoderma reesei*. Preferably, the α -1,2-mannosidase expression vector is
25 engineered such that the α -1,2-mannosidase or a functional part thereof expressed from the vector includes an ER-retention signal. A preferred ER-retention signal is HDEL. The α -1,2-mannosidase coding sequence can be operable linked to a constitutive or inducible promoter, and a 3' termination sequence. The vectors can be

integrative vectors or replicative vectors. Particularly preferred α -1,2-mannosidase expression vectors include pGAPZMFManHDEL, pGAPZMFManMycHDEL, pPICZBMFManMycHDEL, pGAPZmManHDEL, pGAPZmMycManHDEL, pPIC9mMycManHDEL and pGAPZmMycManHDEL.

- 5 In another preferred embodiment, the knock-in vectors of the present invention include a sequence coding for a glucosidase II or a functional part thereof and are capable of expressing the glucosidase II or the functional part in a methylotrophic yeast strain. A preferred nucleotide sequence is a nucleotide sequence encoding the glucosidase II of a fungal species, and more preferably, *Saccharomyces*
10 *cerevisiae*. Preferably, the glucosidase II expression vector is engineered such that the glucosidase II or a functional part thereof expressed from the vector includes an ER-retention signal. A preferred ER-retention signal is HDEL. The glucosidase II coding sequence can be operable linked to a constitutive or inducible promoter, and a 3' termination sequence. The vectors can be integrative vectors or replicative vectors.
- 15 Particularly preferred glucosidase II expression vectors include pGAPZAGLSII, pPICZAGLSII, pAOX2ZAGLSII, pYPTIZAGLSII, pGAPADEglsII, pPICADEglsII, pAOX2ADEglsII, pYPTIADEglsII, pGAPZAglslIIHDEL and pGAPADEglsIIHDEL.

Expression vectors which include both of an α -1,2-mannosidase expression unit and a glucosidase II expression unit are also provided by the present invention.

- 20 In another aspect, the present invention provides "knock-out" vectors which, when introduced into a methylotrophic yeast strain, inactivate or disrupt a gene thereby facilitating the reduction in the glycosylation of glycoproteins produced in the methylotrophic yeast strain.

- In one embodiment, the present invention provides a "knock-out" vector
25 which, when introduced into a methylotrophic yeast strain, inactivates or disrupts the Och1 gene. A preferred Och1 knock-out vector is pBLURA5'PpOCH1.

 Still another embodiment of the present invention provides vectors which include both a knock-in unit and a knock-out unit.

Furthermore, any of the knock-in or knock-out vectors of the present invention can also include a nucleotide sequence capable of expressing a heterologous protein of interest in a methylotrophic yeast.

Another embodiment of the present invention provides methods of
5 modifying the glycosylation in a methylotrophic yeast by transforming the yeast with one or more vectors of the present invention.

Strains of a methylotrophic yeast which can be modified using the present methods include, but are not limited to, yeast strains capable of growth on methanol such as yeasts of the genera *Candida*, *Hansenula*, *Torulopsis*, and *Pichia*. Preferred
10 methylotrophic yeasts are of the genus *Pichia*. Especially preferred are *Pichia pastoris* strains GS115 (NRRL Y-15851), GS190 (NRRL Y-18014), PPF1 (NRRL Y-18017), PPY120H, yGC4, and strains derived therefrom. Methylotrophic yeast strains which can be modified using the present methods also include those methylotrophic yeast strains which have been engineered to express one or more heterologous proteins
15 of interest. The glycosylation on the heterologous proteins expressed from these previously genetically engineered strains can be reduced by transforming such strains with one or more of the vectors of the present invention

Methylotrophic yeast strains which are modified by practicing the present methods are provided in another embodiment of the present invention.

20 A further aspect of the present invention is directed to methods of producing glycoproteins with a reduced glycosylation.

In accordance with such methods, a nucleotide sequence capable of expressing a glycoprotein can be introduced into a methylotrophic yeast strain which has previously been transformed with one or more of the vectors of the present
25 invention. Alternatively, a methylotrophic yeast strain which has been genetically engineered to express a glycoprotein can be transformed with one or more of the vectors of the present invention. Moreover, if a methylotrophic yeast strain is not transformed with a nucleotide sequence encoding a glycoprotein of interest or any of the vectors of the present invention, such yeast strain can be transformed, either

consecutively or simultaneously, with both a nucleotide sequence capable of expressing the glycoprotein and one or more vectors of the present invention.

Additionally, a methylotrophic yeast strain can be transformed with one or more of the present knock-in and/or knock-out vectors which also include a nucleotide sequence

5 capable of expressing a glycoprotein in the methylotrophic yeast strain.

Glycoproteins products produced by using the methods of the present invention, i.e., glycoproteins with reduced N-glycosylation, are also part of the present invention.

Kits which include one or more of the vectors of the present invention, or
10 one or more strains modified to produce glycoproteins with reduced glycosylation, are also provided.

Brief Description of the Drawings

Figure 1 depicts vectors carrying an HDEL-tagged *Trichoderma reesei* α -
15 1,2-mannosidase expression cassette and describes the way in which these vectors were constructed according to methods known in the art. Abbreviations used throughout construction schemes: 5' AOX1 or AOX1 P: *Pichia pastoris* AOX1 promoter sequence; Amp R: ampicillin resistance gene; ColE1: ColE1 origin of replication; 3'AOX1: 3' sequences of the *Pichia pastoris* AOX1 gene; HIS4: HIS4
20 gene of *Pichia pastoris*. AOX TT: transcription terminator sequence of the *Pichia pastoris* AOX1 gene; ORF: open reading frame; S: secretion signal; P TEF1: the promoter sequence of the *Saccharomyces cerevisiae* transcription elongation factor 1 gene; P EM7: synthetic constitutive prokaryotic promoter EM7; Zeocin: Zeocin resistance gene; CYC1 TT: 3' end of the *S. cerevisiae* CYC1 gene; GAP: promoter
25 sequence of the *Pichia pastoris* glyceraldehyde-3-phosphate dehydrogenase gene; PpURA3: *Pichia pastoris* URA3 gene. As can be seen in this figure, the *Trichoderma reesei* α -1,2-mannosidase was operably linked to the coding sequence for the *S. cerevisiae* α -mating factor secretion signal sequence and further operably linked at the 3' terminus of the coding sequence to the coding sequence for an HDEL peptide. The

whole fusion construct was operably linked to either the *P. pastoris* AOX1 promoter (in pPIC9MFManHDEL) or to the *P. pastoris* GAP promoter (in pGAPZMFManHDEL).

Figure 2 depicts vectors carrying an HDEL-tagged *Mus musculus* α -1,2-mannosidase IB expression cassette and describes the way in which these vectors were constructed according to methods known in the art. As can be seen in this figure, the catalytic domain of the *Mus musculus* α -1,2-mannosidase IB was operably linked to the coding sequence for the *S. cerevisiae* α -mating factor secretion signal sequence and further operably linked at the 3' terminus of the coding sequence to the coding sequence for an HDEL peptide. The whole fusion construct was operably linked to either the *P. pastoris* AOX1 promoter (in pPIC9mManHDEL) or to the *P. pastoris* GAP promoter (in pGAPZmManHDEL). Furthermore, variants of the expression cassette were made in which the coding sequence for a cMyc epitope tag was inserted between the coding sequence for the *S. cerevisiae* α -Mating Factor secretion signal sequence and the coding sequence for the catalytic domain of the *Mus musculus* α -1,2-mannosidase IB. This expression cassette was also operably linked to either the *P. pastoris* AOX1 promoter (in pPIC9mMycManHDEL) or to the *P. pastoris* GAP promoter (in pGAPZmMycManHDEL).

Figure 3 depicts vectors carrying a MycHDEL tagged *Trichoderma reesei* α -1,2-mannosidase and the way in which these vectors were obtained. The resulting fusion construction was again operably linked to either the *P. pastoris* AOX1 promoter (in pPICZBMFManMycHDEL) or to the *P. pastoris* GAP promoter (in pGAPZMFManMycHDEL).

Figure 4 demonstrates the intracellular localization of the MycHDEL-tagged *Trichoderma reesei* α -1,2-mannosidase and indicates ER-targeting by immunofluorescence analysis. **Panel A** Western blotting. Yeast strains were grown in 10 ml YPG cultures to an $OD_{600}=10$, diluted fivefold and grown in YPM for 48 h. 1/50th of the culture medium and 1/65th of the cells were analysed by SDS-PAGE and

Western blotting with the mouse monoclonal 9E10 anti-Myc antibody. The position of molecular weight marker proteins are indicated with arrows. Lanes 1-5: cellular lysates. 1,2: pGAPZMFManMycHDEL transformants. 3: non-transformed PPY12OH (negative control). 4,5: pPICZBMFManMycHDEL transformants. Lanes 6-10: culture media. 6: non transformed PPY12OH (negative control). 7,8: pGAPZMFManMycHDEL transformants. 9,10: pPICZBMFManMycHDEL transformants. **Panel B** Immunofluorescence microscopy. 1: phase contrast image of a *P. pastoris* cell (strain PPY12OH transformed with pGAPZMFManHDEL) at 1000x magnification. The nucleus is visible as an ellipse in the lower right quadrant of the cell. 2: same cell as in 1, but in fluorescence microscopy mode to show localization of the *T. reesei* mannosidase-Myc-HDEL protein. The protein is mainly localized in a circular distribution around the nucleus (nuclear envelope), which is typical for an endoplasmic reticulum steady-state distribution. 3: phase contrast image of a *P. pastoris* cell (strain PPY12OH transformed with pGAPZMFManHDEL) at 1000x magnification. 4: same cell in fluorescence microscopy to show localization of the Golgi marker protein OCH1-HA in *P. pastoris* strain PPY12OH. The dot-like distribution throughout the cytoplasm, with 3-4 dots per cell is typical for cis-Golgi distribution in *P. pastoris*.

Figure 5 depicts the co-sedimentation of mannosidase-MycHDEL with Protein Disulfide Isomerase in sucrose density gradient centrifugation. The top panel shows the distribution over the different fractions of the sucrose gradient of the OCH1-HA Golgi marker protein. The middle panel shows this distribution for the Protein Disulfide Isomerase endoplasmic reticulum marker protein. Finally, the bottom panel shows the distribution of the MycHDEL-tagged *Trichoderma reesei* α -1,2-mannosidase over the same fractions. It is concluded that the mannosidase-MycHDEL almost exactly matches the distribution of the ER marker PDI and thus mainly resides in the ER of the *Pichia pastoris* yeast cells.

Figure 6 depicts the N-glycan analysis of *Trypanosoma cruzi* trans-sialidase coexpressed with *Trichoderma reesei* mannosidase-HDEL. Panel A: malto-

oligosaccharide size reference ladder. Sizes of the glycans are expressed in Glucose Units (GU) by comparison of their electrophoretic mobility to the mobility of these malto-oligosaccharides. Panel B: N-glycans derived from recombinant *Trypanosoma cruzi* trans-sialidase expressed in *Pichia pastoris*. The peak at GU=9,2 corresponds to

5 Man₈GlcNAc₂. Panel C: same analytes as panel 2, but after overnight treatment with 3U/ml purified recombinant *T. reesei* α -1,2-mannosidase. Panel D: N-glycans derived from recombinant trans-sialidase co-expressed in *Pichia pastoris* with *T. reesei* mannosidase-HDEL (under control of the GAP promotor). The peak at GU=7,6 corresponds to the Man₅GlcNAc₂ peak in the profile of RNase B (Panel F).

10 Panel E: same analytes as panel D, but after overnight treatment with 3 mU/ml purified recombinant *T. reesei* α -1,2-mannosidase. Panel F: N-glycans derived from bovine RNase B. These glycans consist of Man₅GlcNAc₂ to Man₈GlcNAc₂. Different isomers are resolved, accounting for the number of peaks for Man₇GlcNAc₂.

Figure 7 depicts the processing of influenza haemagglutinin N-glycans by

15 HDEL-tagged *Trichoderma reesei* α -1,2-mannosidase and the HDEL-tagged catalytic domain of murine α -1,2-mannosidase IB. The Man₅GlcNAc₂ reference oligosaccharide runs at scan 1850 in this analysis (not shown). Panel 1: malto-oligosaccharide size reference ladder. Panel 2: N-glycans derived from recombinant influenza haemagglutinin expressed in *Pichia pastoris*. The peak at scan 2250

20 corresponds to Man₉GlcNAc₂. Panel 3: N-glycans derived from recombinant haemagglutinin co-expressed in *Pichia pastoris* with *T. reesei* mannosidase-HDEL (under control of the GAP promotor). The peak at scan 1950 corresponds to Man₆GlcNAc₂. Panel 4: Same analytes as for panel 3, but after overnight treatment with 30 mU purified recombinant *T. reesei* α -1,2-mannosidase. Panel 5: N-glycans

25 derived from recombinant haemagglutinin co-expressed in *Pichia pastoris* with mouse mannosidase IB-HDEL (under control of the GAP promotor). Panel 6: same analytes as for panel 5, but after overnight treatment with 30 mU purified recombinant *T. reesei* α -1,2-mannosidase.

Figure 8 graphically depicts vector pBLURA5'PpOCH1 and the way in which it was constructed.

Figure 9 depicts the scheme for disrupting the *Pichia pastoris* OCH1 gene by single homologous recombination using pBLURA5'PpOCH1.

5 **Figure 10** depicts the cell wall glycoprotein N-glycan analysis of the Och1-inactivated clone and three clones derived from this Och1-inactivated clone by transformation with pGAPZMFManHDEL. Panel 1 shows the analysis of a mixture of malto-oligosaccharides, the degree of polymerisation of which is given by the numbers on the very top of the figure. This analysis serves as a size reference for the
 10 other panels. On the vertical axis of all panels, peak intensity in relative fluorescence units is given. Panel 2-6: analysis of the cell wall glycoprotein N-glycans of the following strains: Panel 2, non-engineered *P. pastoris* strain yGC4; Panel 3, yGC4 transformed with pBLURA5'PpOch1; 4-6, three clones of the strain of Panel 3, supplementarily transformed with pGAPZMFManHDEL. Panel 7: the N-glycans
 15 derived from bovine RNaseB, consisting of a mixture of Man₅₋₉GlcNAc₂. As can be seen from comparison between panel 2 and 3 and reference to panel 7, transformation with pBLURA5'PpOch1 leads to a strongly increased abundance of the Man₈GlcNAc₂ substrate N-glycan (named peak 1 in Panel 2) of OCH1p. Peak 2 represents the Man₉GlcNAc₂ product of OCH1p. Furthermore, upon supplementary transformation
 20 of pGAPZMFManHDEL, the major glycan on the cell wall glycoproteins of three independent clones is the Man₅GlcNAc₂ end product (peak 3 in panel 4) of *T. reesei* α -1,2-mannosidase digestion of the Man₈GlcNAc₂ substrate.

25 **Figure 11** depicts the analysis of exactly the same glycan mixtures as in Figure 10, but after an *in vitro* digest with 3mU/ml purified *Trichoderma reesei* α -1,2-mannosidase, overnight in 20 mM sodium acetate pH=5.0. Axis assignment is the same as in Figure 10. More Man₅GlcNAc₂ is formed in the pBLURA5'PpOch1 transformed strain (Panel 3) than in the parent strain (Panel 2). Peaks in all panels before scan 3900 come from contaminants and should be ignored in the analysis.

Figure 12 depicts the expression vector pGAPZAGLSII (SEQ ID NO: 18).
 P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7:
 synthetic prokaryotic promotor. Zeocin: zeocine resistance marker gene. CYC1 TT:
 transcription terminator of *S. cerevisiae* cytochrome C1 gene. Co1 E1: bacterial origin
 5 of replication. GAP: promotor of the *P. pastoris* GAP gene. GLS2: *S. cerevisiae*
 glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene

Figure 13 depicts the expression vector pAOX2ZAGLSII (SEQ ID NO:
 16). P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7:
 10 synthetic prokaryotic promotor. Zeocin: zeocine resistance marker gene. CYC1 TT:
 transcription terminator of *S. cerevisiae* cytochrome C1 gene. Co1 E1: bacterial origin
 of replication. AOX2 P: promotor of the *P. pastoris* AOX2 gene. GLS2: *S. cerevisiae*
 glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene

Figure 14 depicts the expression vector pPICZAGLSII (SEQ ID NO: 20).
 15 P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7:
 synthetic prokaryotic promotor. Zeocin: zeocine resistance marker gene. CYC1 TT:
 transcription terminator of *S. cerevisiae* cytochrome C1 gene. Co1 E1: origin of
 replication. AOX1 P: promotor of the *P. pastoris* AOX1 gene. GLS2: *S. cerevisiae*
 glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1 gene

Figure 15 depicts the expression vector pYPT1ZAGLSII ((SEQ ID NO:
 22). P TEF1: promotor of *S. cerevisiae* transcription elongation factor gene. P Em7:
 synthetic prokaryotic promotor. Zeocin: zeocine resistance marker gene. CYC1 TT:
 transcription terminator of *S. cerevisiae* cytochrome C1 gene. Co1 E1: origin of
 replication. P YPT1: promotor of the *P. pastoris* YPT1 gene. GLS2: *S. cerevisiae*
 25 glucosidase II gene. AOX1 TT: transcription terminator of the *P. pastoris* AOX1
 gene.

Figure 16 depicts the expression vector pGAPADE1glsII (SEQ ID NO:
 19). Amp R: Ampillicin resistance marker gene. ADE1: *P. pastoris* ADE1 selection

marker gene. GAP: promotor of the P. Pastoris GAP gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene

Figure 17 depicts the expression vector pAOX2ADE1glsII (SEQ ID NO: 17). Amp R: Ampillicin resistance marker gene. ADE1: P. pastoris ADE1 selection
 5 marker gene. AOX2 P: promotor of the P. pastoris AOX2 gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene.

Figure 18 depicts the expression vector pPICADE1glsII (SEQ ID NO: 21). Amp R: Ampillicin resistance marker gene. ADE1: P. pastoris ADE1 selection
 10 marker gene. AOX1 P: promotor of the P. pastoris AOX1 gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene.

Figure 19 depicts the expression vector pYPT1ADE1glsII (SEQ ID NO: 23). Amp R: Ampillicin resistance marker gene. ADE1: P. pastoris ADE1 selection
 15 marker gene. P YPT1: promotor of the P. pastoris YPT1 gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene.

Figure 20 depicts the expression vector pGAPZAglslIHDEL (SEQ ID NO: 24). Amp R: Ampillicin resistance marker gene. ADE1: P. pastoris ADE1 selection
 20 marker gene. GAP: promotor of the P. pastoris GAP gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene.

Figure 21 depicts the expression vector pGAPADE1glsIIHDEL (SEQ ID NO: 25). P TEF1: promotor of S. cerevisiae transcription elongation factor gene. P Em7: synthetic prokaryotic promotor. Zeocin: zeocine resistance marker gene. CYC1 TT: transcription terminator of S. cerevisiae cytochrome C1 gene. Col E1: bacterial origin of replication. GAP: promotor of the P. pastoris GAP gene. GLS2: S. cerevisiae glucosidase II gene. AOX1 TT: transcription terminator of the P. pastoris AOX1 gene.

Figure 22 depicts the test of the GLSII activity assay using a commercially available yeast alpha-glucosidase (Sigma: Cat. No. G-5003). The assay mixture contains phosphate-citrate buffer pH 6.8, mannose, 2-deoxy-D-glucose, the substrate 4-methylumbelliferyl-alpha-D-glucopyranoside and alpha-glucosidase from Sigma. 1: assay mixture illuminated with UV-light after overnight incubation at 37 °C; 2: same as 1, but this time, the assay mixture lacks the alpha-glucosidase; 3: same as 1, but this time, the assay mixture lacks the substrate.

Figure 23 depicts the results of the activity of recombinantly expressed GLSII from *Pichia pastoris*. All assay mixtures were incubated overnight at 37 °C and afterwards illuminated with UV-light. 1: assay with yeast alpha-glucosidase (Sigma: Cat. No. G-5003); 2: assay with the purified medium of strain 18 (PPY12-OH transformed with pGAPZAGLSII); 3: assay with purified medium of the WT PPY12-OH strain; 4: assay with the purified medium of strain H3 (PPY12-OH transformed with pGAPZAglslIHDEL).

15

Detailed Description of the Invention

It has been established that the majority of N-glycans on glycoproteins leaving the endoplasmic reticulum (ER) of *Pichia* have the core Man₈GlcNAc₂ oligosaccharide structure. After the proteins are transported from the ER to the Golgi apparatus, additional mannose residues are added to this core sugar moiety by different mannosyltransferases, resulting in glycoproteins with large mannose side chains. Such hyperglycosylation of recombinant glycoproteins is undesirable in many instances. Accordingly, the present invention provides methods and vectors for genetically modifying methylotrophic yeast strains to produce glycoproteins with reduced glycosylation. Methylotrophic yeast strains generated using the present methods and vectors, as well as glycoproteins produced from such genetically modified strains are also provided.

In one embodiment, the present invention provides vectors useful for genetically modifying methylotrophic yeast strains to produce glycoproteins with reduced glycosylation.

In one aspect, the present invention provides "knock-in" vectors which are
5 capable of expressing in a methylotrophic yeast strain one or more proteins whose enzymatic activities lead to a reduction of glycosylation in glycoproteins produced by the methylotrophic yeast strain. According to the present invention, such proteins include, e.g., an α -1,2-mannosidase, a glucosidase II, or functional parts thereof.

In a preferred embodiment, the vectors of the present invention include a
10 sequence coding for an α -1,2-mannosidase or a functional part thereof and are capable of expressing the α -1,2-mannosidase or the functional part in a methylotrophic yeast strain.

An α -1,2-mannosidase cleaves the α -1,2-linked mannose residues at the non-reducing ends of $\text{Man}_8\text{GlcNAc}_2$, and converts this core oligosaccharide on
15 glycoproteins to $\text{Man}_5\text{GlcNAc}_2$. *In vitro*, $\text{Man}_5\text{GlcNAc}_2$ is a very poor substrate for any *Pichia* Golgi mannosyltransferase, i.e., mannose residues can not be added to this sugar structure. On the other hand, $\text{Man}_5\text{GlcNAc}_2$ is the acceptor substrate for the mammalian N-acetylglucosaminyl-transferase I and is an intermediate for the hybrid- and complex-type sugar chains characteristic of mammalian glycoproteins. Thus, by
20 way of introducing an α -1,2-mannosidase into methylotrophic yeasts such as *Pichia*, glycoproteins with reduced mannose content can be produced.

According to the present invention, the nucleotide sequence encoding an α -1,2-mannosidase for use in the expression vector of the present invention can derive from any species. A number of α -1,2-mannosidase genes have been cloned and are
25 available to those skilled in the art, including mammalian genes encoding, e.g., a murine α -1,2-mannosidase (Herscovics et al. *J. Biol. Chem.* 269: 9864-9871, 1994), a rabbit α -1,2-mannosidase (Lal et al. *J. Biol. Chem.* 269: 9872-9881, 1994) or a human α -1,2-mannosidase (Tremblay et al. *Glycobiology* 8: 585-595, 1998), as well as fungal

genes encoding, e.g., an *Aspergillus* α -1,2-mannosidase (msdS gene), a *Trichoderma reesei* α -1,2-mannosidase (Maras et al. *J. Biotechnol.* 77: 255-263, 2000), or a *Saccharomyces cerevisiae* α -1,2-mannosidase. Protein sequence analysis has revealed a high degree of conservation among the eukaryotic α -1,2-mannosidases identified so far.

Preferably, the nucleotide sequence for use in the present vectors encodes a fungal α -1,2-mannosidase, more preferably, a *Trichoderma reesei* α -1,2-mannosidase, and more particularly, the *Trichoderma reesei* α -1,2-mannosidase described by Maras et al. *J. Biotechnol.* 77: 255-63 (2000).

According to the present invention, the nucleotide sequence can also code for only a functional part of an α -1,2-mannosidase.

By "functional part" is meant a polypeptide fragment of an α -1,2-mannosidase which substantially retains the enzymatic activity of the full-length protein. By "substantially" is meant at least about 40%, or preferably, at least 50% or more of the enzymatic activity of the full-length α -1,2-mannosidase is retained. For example, as illustrated by the present invention, the catalytic domain of the murine α -1,2-mannosidase IB constitutes a "functional part" of the murine α -1,2-mannosidase IB. Those skilled in the art can readily identify and make functional parts of an α -1,2-mannosidase using a combination of techniques known in the art. Predictions of the portions of an α -1,2-mannosidase essential to or sufficient to confer the enzymatic activity can be made based on analysis of the protein sequence. The activity of a portion of an α -1,2-mannosidase of interest, expressed and purified from an appropriate expression system, can be verified using *in vitro* or *in vivo* assays described hereinbelow.

In accordance with the present invention, an α -1,2-mannosidase or a functional part thereof expressed in a methylotrophic yeast strain preferably is targeted to a site in the secretory pathway where $\text{Man}_8\text{GlcNAc}_2$ (the substrate of α -1,2-

mannosidase) is already formed on a glycoprotein, but has not reached a Golgi glycosyltransferase which elongates the sugar chain with additional mannose residues.

Accordingly, in a preferred embodiment of the present invention, the α -1,2-mannosidase expression vector is engineered as such that the α -1,2-mannosidase
5 or a functional part thereof expressed from the vector includes an ER-retention signal.

"An ER retention signal" refers to a peptide sequence which directs a protein having such peptide sequence to be transported to and retained in the ER. Such ER retention sequences are often found in proteins that reside and function in the ER.

10 Multiple choices of ER retention signals are available to those skilled in the art, e.g., the first 21 amino acid residues of the *S. cerevisiae* ER protein MNS1 (Martinet et al. *Biotechnology Letters* 20: 1171-1177, 1998). A preferred ER retention signal for use in the present invention is peptide HDEL (SEQ ID NO: 1). The HDEL peptide sequence, found in the C-terminus of a number of yeast proteins, acts as a
15 retention/retrieval signal for the ER (Pelham *EMBO J.* 7: 913-918, 1988). Proteins with an HDEL sequence are bound by a membrane-bound receptor (Erd2p) and then enter a retrograde transport pathway for return to the ER from the Golgi apparatus.

According to the present invention, an ER retention signal can be placed anywhere in the protein sequence of an α -1,2-mannosidase, but preferably at the C-
20 terminus of the α -1,2-mannosidase.

The α -1,2-mannosidase for use in the present invention can be further modified, e.g., by insertion of an epitope tag to which antibodies are available, such as Myc, HA, FLAG and His6 tags well-known in the art. An epitope-tagged α -1,2-mannosidase can be conveniently purified, or monitored for both expression and
25 intracellular localization.

An ER retention signal and an epitope tag can be readily introduced into a protein of interest by inserting nucleotide sequences coding for such signal or tag into

the nucleotide sequence encoding the protein of interest, using any of the molecular biology techniques known in the art.

In another preferred embodiment, the vectors of the present invention include a sequence coding for a glucosidase II or a functional part thereof and are
5 capable of expressing the glucosidase II or the functional part in the methylotrophic yeast strain.

It has been established that the initial N-linked oligosaccharide (Glc₃Man₉GlcNAc₂), transferred in the ER onto a protein, is cleaved in the ER by specific glucosidases to remove the glucose residues, and by a mannosidase to remove
10 one specific α -1,2-linked mannose. It has been observed by the present inventors that some recombinant proteins expressed in *Pichia* have residual glucose residues on the sugar moiety when such proteins leave the ER for the Golgi apparatus. The residual glucose molecules present on the sugar structure prevent the complete digestion of the sugar moiety by an α -1,2-mannosidase, and the introduction of an exogenous
15 glucosidase can facilitate the removal of these glucose residues.

According to the present invention, the nucleotide sequence encoding a glucosidase II can derive from any species. Glucosidase II genes have been cloned from a number of mammalian species including rat, mouse, pig and human. The glucosidase II protein from these mammalian species consists of an alpha and a beta
20 subunit. The alpha subunit is about 110 kDa and contains the catalytic activity of the enzyme, while the beta subunit has a C-terminal HDEL ER-retention sequence and is believed to be important for the ER localization of the enzyme. The glucosidase II gene from *S. cerevisiae* has also been cloned (ORF YBR229c, located on chromosome II). This gene encodes a protein of about 110 kDa, which shows a high degree of
25 homology to the mammalian alpha subunits.

A preferred glucosidase II gene for use in the present vectors is from a fungal species such as *Pichia pastoris* and *S. cerevisiae*. An example of a fungal glucosidase II gene is the *S. cerevisiae* glucosidase II alpha subunit gene.

According to the present invention, the nucleotide sequence can also encode only a functional part of a glucosidase II. By "functional part" is meant a polypeptide fragment of a glucosidase II which substantially retains the enzymatic activity of the full-length protein. By "substantially" is meant at least about 40%, or
5 preferably, at least 50% or more of the enzymatic activity of the full-length glucosidase II is retained. Functional parts of a glucosidase II can be identified and made by those skilled in the art using a variety of techniques known in the art.

In a preferred embodiment of the present invention, the glucosidase II protein is engineered to include an ER retention signal such that the protein expressed
10 in a methylotrophic yeast strain is targeted to the ER and retains therein for function. ER retention signals are as described hereinabove, e.g., the HDEL peptide sequence.

The glucosidase II for use in the present invention can be further modified, e.g., by insertion of an epitope tag to which antibodies are available, such as Myc, HA, FLAG, and His6 tag, which are well-known in the art.

15 According to the present invention, the "knock-in" vectors can include either or both of an α -1,2-mannosidase coding sequence and a glucosidase II coding sequence.

Further according to the present invention, the nucleotide sequence coding for the enzyme to be expressed (e.g., an α -1,2-mannosidase or a functional part
20 thereof, or a glucosidase II or a functional part thereof) can be placed in an operable linkage to a promoter and a 3' termination sequence.

Promoters appropriate for expression of a protein in a methylotrophic yeast can include both constitutive promoters and inducible promoters. Constitutive promoters include e.g., the *Pichia pastoris* glyceraldehyde-3-phosphate
25 dehydrogenase promoter ("the GAP promoter"). Examples of inducible promoters include, e.g., the *Pichia pastoris* alcohol oxidase I promoter ("the AOXI promoter") (U.S. Patent No. 4,855,231), or the *Pichia pastoris* formaldehyde dehydrogenase promoter ("the FLD promoter") (Shen et al. Gene 216: 93-102, 1998).

3' termination sequences are sequences 3' to the stop codon of a structural gene which function to stabilize the mRNA transcription product of the gene to which the sequence is operably linked, such as sequences which elicit polyadenylation. 3' termination sequences can be obtained from *Pichia* or other methylotrophic yeast.

- 5 Examples of *Pichia pastoris* 3' termination sequences useful for the practice of the present invention include termination sequences from the *AOX1* gene, *p40* gene, *HIS4* gene and *FLD1* gene.

The vectors of the present invention preferably contain a selectable marker gene. The selectable marker may be any gene which confers a selectable phenotype
10 upon a methylotrophic yeast strain and allows transformed cells to be identified and selected from untransformed cells. The selectable marker system may include an auxotrophic mutant methylotrophic yeast strain and a wild type gene which complements the host's defect. Examples of such systems include the *Saccharomyces cerevisiae* or *Pichia pastoris HIS4* gene which may be used to complement *his4*
15 *Pichia* strains, or the *S. cerevisiae* or *Pichia pastoris ARG4* gene which may be used to complement *Pichia pastoris arg* mutants. Other selectable marker genes which function in *Pichia pastoris* include the *Zeo^R* gene, the *G418^R* gene, and the like.

The vectors of the present invention can also include an autonomous replication sequence (ARS). For example, U.S. Patent No. 4,837,148 describes
20 autonomous replication sequences which provide a suitable means for maintaining plasmids in *Pichia pastoris*. The disclosure of U.S. Patent No. 4,837,148 is incorporated herein by reference.

The vectors can also contain selectable marker genes which function in bacteria, as well as sequences responsible for replication and extrachromosomal
25 maintenance in bacteria. Examples of bacterial selectable marker genes include ampicillin resistance (*Amp^r*), tetracycline resistance (*Tet^r*), neomycin resistance, hygromycin resistance, and zeocin resistance (*Zeo^R*) genes.

According to the present invention, the nucleotide sequence encoding the protein to be expressed in a methylotrophic yeast can be placed in an integrative vector or a replicative vector (such as a replicating circular plasmid).

Integrative vectors are disclosed, e.g., in U.S. Patent No. 4,882,279 which is incorporated herein by reference. Integrative vectors generally include a serially arranged sequence of at least a first insertable DNA fragment, a selectable marker gene, and a second insertable DNA fragment. The first and second insertable DNA fragments are each about 200 nucleotides in length and have nucleotide sequences which are homologous to portions of the genomic DNA of the species to be transformed. A nucleotide sequence containing a structural gene of interest for expression is inserted in this vector between the first and second insertable DNA fragments whether before or after the marker gene. Integrative vectors can be linearized prior to yeast transformation to facilitate the integration of the nucleotide sequence of interest into the host cell genome.

Replicative and integrative vectors carrying either or both of an α -1,2-mannosidase coding sequence or a glucosidase II coding sequence can be constructed by standard techniques known to one of ordinary skill in the art and found, for example, in Sambrook et al. (1989) in *Molecular Cloning: A Laboratory Manual*, or any of a myriad of laboratory manuals on recombinant DNA technology that are widely available.

Preferred vectors of the present invention carrying an α -1,2-mannosidase expression sequence include pGAPZMFManHDEL, pGAPZMFManMycHDEL, pPICZBMFManMycHDEL, pGAPZmManHDEL, pGAPZmMycManHDEL, pPIC9mMycManHDEL and pGAPZmMycManHDEL, which are further described in the Examples hereinbelow.

Preferred vectors of the present invention carrying a glucosidase II expression sequence include pGAPZAGLSII, pPICZAGLSII, pAOX2ZAGLSII, pYPTIZAGLSII, pGAPADE1glsII, pPICADE1glsII, pAOX2ADE1glsII,

pYPTIADE1glsII, pGAPZAglslIHDEL and pGAPADE1glsIIHDEL, which are further described in the Examples hereinbelow.

In another aspect, the present invention provides "knock-out" vectors which, when introduced into a methylotrophic yeast strain, inactivate or disrupt a gene thereby facilitating the reduction in the glycosylation of glycoproteins produced in the methylotrophic yeast strain.

In one embodiment, the present invention provides a "knock-out" vector which, when introduced into a methylotrophic yeast strain, inactivates or disrupts the Och1 gene.

The *S. cerevisiae* OCH1 gene has been cloned (Nakayama et al. *EMBO J.* 11: 2511-2519, 1992). It encodes a membrane bound α -1,6-mannosyltransferase, localized in the early Golgi complex, that is functional in the initiation of α -1,6-polymannose outer chain addition to the N-linked core oligosaccharide (Man₅GlcNAc₂ and Man₈GlcNAc₂) (Nakanishi-Shindo et al. *J. Biol. Chem.* 268: 26338-26345, 1993).

A *Pichia* sequence has been described in Japanese Patent Application No. 07145005 that encodes a protein highly homologous to the *S. cerevisiae* OCH1. For purpose of the present invention, this sequence is denoted herein as "the *Pichia* OCH1 gene". Those skilled in the art can isolate the OCH1 genes from other methylotrophic yeasts using techniques well known in the art.

According to the present invention, a disruption in the OCH1 gene of a methylotrophic yeast can result in either the production of an inactive protein product or no product. The disruption may take the form of an insertion of a heterologous DNA sequence into the coding sequence and/or the deletion of some or all of the coding sequence. Gene disruptions can be generated by homologous recombination essentially as described by Rothstein (in *Methods in Enzymology*, Wu et al., eds., vol 101:202-211, 1983).

To disrupt the Och1 gene by homologous recombination, an Och1 knock-out vector can be constructed in such a way to include a selectable marker gene. The selectable marker gene is operably linked, at both 5' and 3' end, to portions of the

Och1 gene of sufficient length to mediate homologous recombination. The selectable marker can be one of any number of genes which either complement host cell auxotrophy or provide antibiotic resistance, including *URA3*, *LEU2* and *HIS3* genes. Other suitable selectable markers include the *CAT* gene, which confers
5 chloramphenicol resistance on yeast cells, or the *lacZ* gene, which results in blue colonies due to the expression of active β -galactosidase. Linearized DNA fragments of an Och1 knock-out vector are then introduced into host methylotrophic yeast cells using methods well known in the art. Integration of the linear fragments into the genome and the disruption of the Och1 gene can be determined based on the selection
10 marker and can be verified by, for example, Southern Blot analysis.

Alternatively, an Och1 knock-out vector can be constructed in such a way to include a portion of the Och1 gene to be disrupted, which portion is devoid of any Och1 promoter sequence and encodes none or an inactive fragment of the Och1 protein. By "an inactive fragment", it is meant a fragment of the Och1 protein which
15 has, preferably, less than about 10% and most preferably, about 0% of the activity of the full-length OCH1 protein. Such portion of the OCH1 gene is inserted in a vector in such a way that no known promoter sequence is operably linked to the OCH1 sequence, but that a stop codon and a transcription termination sequence are operably linked to the portion of the Och1 gene. This vector can be subsequently linearized in
20 the portion of the OCH1 sequence and transformed into a methylotrophic yeast strain using any of the methods known in the art. By way of single homologous recombination, this linearized vector is then integrated in the OCH1 gene. Two Och1 sequences are produced in the chromosome as a result of the single homologous recombination. The first Och1 sequence is the portion of the Och1 gene from the
25 vector, which is now under control of the OCH1 promoter of the host methylotrophic yeast, yet cannot produce an active OCH1 protein as such Och1 sequence codes for no or an inactive fragment of the OCH1 protein, as described hereinabove. The second Och1 sequence is a full OCH1 coding sequence, but is not operably linked to any known promoter sequence and thus, no active messenger is expected to be formed for

synthesis of an active OCH1 protein. Preferably, an inactivating mutation is introduced in the OCH1 sequence, to the 5' end of the site of linearization of the vector and to the 3' end of the translation initiation codon of OCH1. By "inactivating mutation" it is meant a mutation introducing a stop codon, a frameshift mutation or
5 any other mutation causing a disruption of the reading frame. Such mutation can be introduced into an Och1 sequence using any of the site directed mutagenesis methods known in the art. Such inactivating mutation ensures that no functional OCH1 protein can be formed even if there exist some promoter sequences 5' to the Och1 sequence in the knock-out vector.

10 A preferred Och1 knock-out vector of the present invention is pBLURA5'PpOCH1.

If desired, either or both of a mannosidase expression sequence and a glucosidase expression sequence can be carried on the same plasmid used to disrupt the OCH1 gene to create a "knock-in-and-knock-out" vector.

15 Additionally, any of the above-described vectors can further include a nucleotide sequence capable of expressing a glycoprotein of interest in a methylotrophic yeast strain.

Another aspect of the present invention is directed to methods of modifying methylotrophic yeast strains to reduce glycosylation on proteins produced by the
20 methylotrophic yeast strains. In accordance with the present methods, methylotrophic yeast strains are modified by transforming into these yeast strains with one or more, i.e., at least one, knock-in and/or knock-out vectors of the present invention as described herein above.

Methylotrophic yeast strains which can be modified using the present
25 methods include but are not limited to yeast capable of growth on methanol such as yeasts of the genera *Candida*, *Hansenula*, *Torulopsis*, and *Pichia*. A list of species which are exemplary of this class of yeasts can be found in C. Anthony (1982), *The Biochemistry of Methylotrophs*, 269. *Pichia pastoris*, *Pichia methanolica*, *Pichia anomala*, *Hansenula polymorpha* and *Candida boidinii* are examples of

methyiotrophic yeasts useful in the practice of the present invention. Preferred methyiotrophic yeasts are of the genus *Pichia*. Especially preferred are *Pichia pastoris* strains GS115 (NRRL Y-15851); GS190 (NRRL Y-18014) disclosed in U.S. Patent No. 4,818,700; PPF1 (NRRL Y-18017) disclosed in U.S. Patent No. 4,812,405; 5 PPY120H and yGC4; as well as strains derived therefrom.

Methyiotrophic yeast strains which can be modified using the present methods also include those methyiotrophic yeast strains which have been genetically engineered to express one or more heterologous glycoproteins of interest. The glycosylation on the heterologous glycoproteins expressed from these previously 10 engineered strains can be reduced by transforming such strains with one or more of the vectors of the present invention.

The vectors of the present invention can be introduced into the cells of a methyiotrophic yeast strain using known methods such as the spheroplast technique, described by Cregg et al. 1985, or the whole-cell lithium chloride yeast transformation 15 system, Ito et al. *Agric. Biol. Chem.* 48:341, modified for use in *Pichia* as described in EP 312,934. Other published methods useful for transformation of the plasmids or linear vectors include U.S. Patent No. 4,929,555; Hinnen et al. *Proc. Nat. Acad. Sci. USA* 75:1929 (1978); Ito et al. *J. Bacteriol.* 153:163 (1983); U.S. Patent No. 4,879,231; Sreekrishna et al. *Gene* 59:115 (1987). Electroporation and PEG1000 20 whole cell transformation procedures may also be used. Cregg and Russel *Methods in Molecular Biology: Pichia Protocols*, Chapter 3, Humana Press, Totowa, N.J., pp. 27-39 (1998).

Transformed yeast cells can be selected by using appropriate techniques including but not limited to culturing auxotrophic cells after transformation in the 25 absence of the biochemical product required (due to the cell's auxotrophy), selection for and detection of a new phenotype, or culturing in the presence of an antibiotic which is toxic to the yeast in the absence of a resistance gene contained in the transformants. Transformants can also be selected and/or verified by integration of the

expression cassette into the genome, which can be assessed by e.g., Southern Blot or PCR analysis.

In one embodiment, a methylotrophic yeast strain is transformed with a vector which includes a nucleotide sequence coding for an α -1,2-mannosidase or a functional part thereof. The nucleotide sequence is capable of expressing the α -1,2-mannosidase or the functional part in the methylotrophic yeast strain, and is, preferably, integrated into the genome of the methylotrophic yeast strain.

The expression of an α -1,2-mannosidase introduced in a methylotrophic yeast strain can be verified both at the mRNA level, e.g., by Northern Blot analysis, and at the protein level, e.g., by Western Blot analysis. The intracellular localization of the protein can be analyzed by using a variety of techniques, including subcellular fractionation and immunofluorescence experiments. An ER localization of an α -1,2-mannosidase can be determined by co-sedimentation of this enzyme with a known ER resident protein (e.g., Protein Disulfide Isomerase) in a subcellular fractionation experiment. An ER localization can also be determined by an immunofluorescence staining pattern characteristic of ER resident proteins, typically a perinuclear staining pattern.

To confirm that an α -1,2-mannosidase or a functional part thereof expressed in a methylotrophic yeast strain has the expected mannose-trimming activity, both *in vitro* and *in vivo* assays can be employed. Typically, an *in vitro* assay involves digestion of an *in vitro* synthesized substrate, e.g., $\text{Man}_8\text{GlcNAc}_2$, with the enzyme expressed and purified from a methylotrophic yeast strain, and assessing the ability of such enzyme to trim $\text{Man}_8\text{GlcNAc}_2$ to, e.g., $\text{Man}_5\text{GlcNAc}_2$. In *in vivo* assays, the α -1,2-mannosidase or a part thereof is co-expressed in a methylotrophic yeast with a glycoprotein known to be glycosylated with N-glycans bearing terminal α -1,2-linked mannose residues in such yeast. The enzymatic activity of such an α -1,2-mannosidase or a part thereof can be measured based on the reduction of the number of α -1,2-linked mannose residues in the structures of the N-glycans of the

glycoprotein. In both *in vitro* and *in vivo* assays, the composition of a carbohydrate group can be determined using techniques that are well known in the art and are illustrated in the Examples hereinbelow.

In another embodiment, a methylotrophic yeast strain is transformed with a
5 vector which includes a nucleotide sequence coding for a glucosidase II or a functional part thereof. The nucleotide sequence is capable of expressing the glucosidase II or the functional part in the methylotrophic yeast strain, and is, preferably, integrated into the genome of the methylotrophic yeast strain.

The enzymatic activity of a glucosidase II or a functional part thereof
10 expressed in a transformed methylotrophic yeast strain can be assessed using a variety of assays. For example, methylotrophic yeast cells transformed with a sequence encoding a glucosidase II or a part thereof can be set to grow on solid medium containing a substrate of the glucosidase, e.g., 5-bromo-4-chloro-3-indolyl- α -D-glucopyranoside or 4-MU- α -D-Glc. When the enzyme is expressed by the *Pichia* and
15 secreted extracellularly, the substrate is acted upon by the enzyme, giving rise to detectable signals around the colonies such as blue color or fluorescent glow. Alternatively, liquid culture medium containing the expressed protein molecules can be collected and incubated in test tubes with a substrate, e.g., p-nitrophenyl- α -D-glucopyranoside. The enzymatic activity can be determined by measuring the specific
20 product released. Moreover, *in vivo* assays can be employed, where a glucosidase II is co-expressed in yeast with a glycoprotein known to be N-glycosylated with glucose residues, e.g., influenza neuraminidase. The enzymatic activity of the glucosidase II can be measured based on the reduction of the glucose content in the sugar chain(s) of the glycoprotein.

25 In still another embodiment of the present invention, a methylotrophic yeast strain is transformed with an Och1 knock-out vector. As a result of the transformation and integration of the vector, the genomic Och1 gene in the yeast strains is disrupted.

In a further embodiment of the present invention, a methylotrophic yeast strain is transformed with any combination of an α -1,2-mannosidase expression vector, a glucosidase II expression vector, and an Och1 knock-out vector. Such modification can be achieved by serial, consecutive transformations, i.e., introducing
5 one vector at a time, or alternatively by co-transformation, i.e., introducing the vectors simultaneously.

The modified methylotrophic yeast strains described herein above can be further modified if desired. For example, additional disruption of genes encoding any other *Pichia* mannosyltransferases can be made. Genes encoding mammalian
10 enzymes can also be introduced to produce glycoproteins having hybrid- or complex-type N-glycans, if desired.

Methylotrophic yeast strains which are modified by using the present methods, i.e., by transforming with one or more of the vectors of the present invention, form another embodiment of the present invention.

15 It should be understood that certain aspects of the present invention, especially the introduction of an intracellularly expressed α -1,2-mannosidase activity, are also useful to obtain a reduced glycosylation of the O-linked glycans on glycoproteins produced in a methylotrophic yeast, as it is known in the art that these O-linked glycans consist mainly of α -1,2-linked mannose residues. O-linked glycans
20 as used herein refers to carbohydrate structures linked to serine or threonine residues of glycoproteins.

A further aspect of the invention is directed to methods of producing a glycoprotein with reduced glycosylation in a methylotrophic yeast, especially a glycoprotein heterologous to the methylotrophic yeast.

25 "A glycoprotein" as used herein refers to a protein which, in methylotrophic yeasts, is either glycosylated on one or more asparagines residues or on one or more serine or threonine residues, or on both asparagines and serine or threonine residues.

The term "reduced glycosylation" refers to a reduced size of the carbohydrate moiety on the glycoprotein, particularly with fewer mannose residues, when the glycoprotein is expressed in a methylotrophic yeast strain which has been modified in accordance with the present invention, as compared to a wild type, unmodified strain of the methylotrophic yeast.

In accordance with the present invention, the production of a glycoprotein of interest with reduced glycosylation can be achieved in a number of ways. A nucleotide sequence capable of expressing a glycoprotein can be introduced into a methylotrophic yeast strain which has been previously modified in accordance with the present invention, i.e., a strain transformed with one or more of the vectors of the present invention and capable of producing glycoproteins with reduced glycosylation. Alternatively, a methylotrophic yeast strain which has already been genetically engineered to express a glycoprotein can be transformed with one or more of the vectors of the present invention. Otherwise, if a methylotrophic yeast strain does not express a glycoprotein of interest, nor is the strain transformed with any of the vectors of the present invention, such yeast strain can be transformed, either consecutively or simultaneously, with both a nucleotide sequence capable of expressing the glycoprotein and one or more vectors of the present invention. Additionally, a methylotrophic yeast strain can be transformed with one or more of the present knock-in and/or knock-out vectors which also include a nucleotide sequence capable of expressing a glycoprotein in the methylotrophic yeast strain.

The nucleotide sequence capable of expressing a glycoprotein in a methylotrophic yeast can be made to include from 5' to 3', a promoter, a sequence encoding the glycoprotein, and a 3' termination sequence. Promoters and 3' termination sequences which are suitable for expression of a glycoprotein can include any of those promoters and 3' termination sequences described hereinabove.

The nucleotide sequence for expression of a glycoprotein can include additional sequences, e.g., signal sequences coding for transit peptides when secretion of a protein product is desired. Such sequences are widely known, readily available

and include *Saccharomyces cerevisiae* alpha mating factor prepro (α mf), *Pichia pastoris* acid phosphatase (PHO1) signal sequence and the like.

The nucleotide sequence for expression of a glycoprotein can be placed on a replicative vector or an integrative vector. The choice and construction of such
5 vectors are as described hereinabove.

The nucleotide sequence capable of expressing a glycoprotein can be carried on the same replicative plasmid as a plasmid-borne α -1,2-mannosidase or glucosidase II expression unit. Alternatively, the nucleotide sequence containing the glycoprotein coding sequence is carried on a separate plasmid or integrated into the
10 host genome.

Glycoproteins produced can be purified by conventional methods. Purification protocols can be determined by the nature of the specific protein to be purified. Such determination is within the ordinary level of skill in the art. For example, the cell culture medium is separated from the cells and the protein secreted
15 from the cells can be isolated from the medium by routine isolation techniques such as precipitation, immunoadsorption, fractionation or a variety of chromatographic methods.

Glycoproteins which can be produced by the methods of the present invention include, e.g., *Bacillus amyloliquefaciens* α -amylase, *S. cerevisiae* invertase,
20 *Trypanosoma cruzi* trans-sialidase, HIV envelope protein, influenza virus A haemagglutinin, influenza neuraminidase, Bovine herpes virus type-1 glycoprotein D, human angiostatin, human B7-1, B7-2 and B-7 receptor CTLA-4, human tissue factor, growth factors (e.g., platelet-derived growth factor), tissue plasminogen activator, plasminogen activator inhibitor-I, urokinase, human lysosomal proteins such as α -
25 galactosidase, plasminogen, thrombin, factor XIII and immunoglobulins. For additional useful glycoproteins which can be expressed in the genetically engineered *Pichia* strains of the present invention, see Bretthauer and Castellino, *Biotechnol.*

Appl. Biochem. 30: 193-200 (1999), and Kukuruzinska et al. *Ann Rev. Biochem.* 56: 915-44 (1987).

Glycoproteins produced by using the methods of the present invention, i.e., glycoproteins with reduced glycosylation, are also part of the present invention.

5 Still another aspect of the present invention provides kits which contain one or more of the knock-in vectors, knock-out vectors, or knock-in-and-knock-out vectors of the present invention described above. More particularly, a kit of the present invention contains a vector capable of expressing an α -mannosidase I in a methylotrophic yeast, a vector capable of expressing a glucosidase II in a
10 methylotrophic yeast, a vector capable of disrupting the Och1 gene in a methylotrophic yeast, a vector capable of expressing both a glucosidase II and an α -mannosidase, a vector a vector capable of disrupting the Och1 gene and capable of expressing either or both of a glucosidase II and an α -mannosidase, or any combinations thereof.

15 The kit can also include a nucleotide sequence which encodes and is capable of expressing a heterologous glycoprotein of interest. Such nucleotide sequence can be provided in a separate vector or in the same vector which contains sequences for knocking-in or knocking out as described hereinabove.

20 In addition, the kit can include a plasmid vector in which a nucleotide sequence encoding a heterologous protein of interest can be subsequently inserted for transformation into and expression in a methylotrophic yeast. Alternatively, the knock-in or knock-out vectors in the kits have convenient cloning sites for insertion of a nucleotide sequence encoding a heterologous protein of interest.

25 The kit can also include a methylotrophic yeast strain which can be subsequently transformed with any of the knock-in, knock-out or knock-in-and-knock-out vectors described hereinabove. The kit can also include a methylotrophic yeast strain which has been transformed with one or more of the knock-in or knock-out vectors. Furthermore, the kit can include a methylotrophic yeast strain which has been

transformed with a nucleotide sequence encoding and capable of expressing a heterologous glycoprotein of interest.

The present invention is further illustrated by the following examples.

Example 1**Introduction of α -1,2-Mannosidase to the ER-Golgi Border****1.1 Plasmids**

Plasmid	Promoter	Enzyme	Tag
pGAPZMFManHDEL	GAP	<i>T. reesei</i> α -1,2-mannosidase	--
pGAPZMFManMycHDEL	GAP	<i>T. reesei</i> α -1,2-mannosidase	Myc
pPICZBMFManMycHDEL	AOX1	<i>T. reesei</i> α -1,2-mannosidase	Myc
pGAPZMFmManHDEL	GAP	mouse mannosidase IB catalytic domain	--
pGAPZMFmMycManHDEL	GAP	mouse mannosidase IB catalytic domain	Myc

5

The *Trichoderma reesei* α -1,2-mannosidase gene has been isolated and described by Maras et al. (*J. Biotechnol.* 77;255-263, 2000). The sequence of this gene is available at NCBI Genbank under Accession No. AF212153. A construction fragment was generated by PCR using the pPIC9MFmanase plasmid (same as pPP1MFmds1 described by Maras et al. (2000)) as the template and using the following oligonucleotide primers: 5'-GACTGGTTCCAATTGACAAGC-3' (SEQ ID NO:2) and 5'-AGTCTAGATTACAACTCGTCGTGAGCAAGGTGGCCGCCCCGTCG-3' (SEQ ID NO:3). The resulting product contained the 3' end of the *Pichia pastoris* AOX1 promoter, the prepro-signal sequence of the *S. cerevisiae* α -mating factor, the open reading frame of the *Trichoderma reesei* α -1,2-mannosidase cloned in frame with the signal sequence, the coding sequence for HDEL, a stop codon and an *Xba* I restriction site. This fragment was digested with *Eco* RI and *Xba* I, removing the 5' sequences up to the mannosidase ORF, and then cloned into the vector pGAPZ α A (Invitrogen, Baarn, The Netherlands) which had been digested with *Eco* RI and *Xba* I, thus restoring the fusion with the *S. cerevisiae* α -mating factor signal

20

sequence. The resulting plasmid was named pGAPZMFMHDEL and is graphically depicted in **Figure 1**. The ORF sequence of the MFMHDEL fusion in pGAPZMFMHDEL is set forth in SEQ ID NO: 14.

In order to introduce the coding sequence for a c-Myc tag between the
 5 catalytic domain and the HDEL-signal, the 3' end of the ORF of *T. reesei* α -1,2-mannosidase was PCR-amplified using a sense primer 5'-CCATTGAGGACGCATGCCGCGCC-3' (SEQ ID NO: 4) (containing an *Sph* I restriction site) and an antisense primer
 GTATCTAGATTACAACTCGTCGTGCAGATCCTCTTCTGAGATGAGTTTTTGT
 10 TCAGCAAGGTGGCCGCCCCGTCGTGATGATGAA (SEQ ID NO: 5) (containing the coding sequences of the c-Myc tag and the HDEL signal, followed by a stop codon and an *Xba* I restriction site). The resulting PCR product was digested with *Sph* I and *Xba* I, purified by agarose gel electrophoresis and inserted into pGAPZMFMHDEL which had been cut with the same restriction enzymes, resulting in plasmid
 15 pGAPZMFMHMyHDEL. To put the ORF of pGAPZMFMHMyHDEL under the control of the inducible AOXI promoter, the entire ORF was liberated from pGAPZMFMHMyHDEL with *Bst* BI and *Xba* I, and cloned in pPICZB (Invitrogen, Baarn, The Netherlands), resulting in pPICZBMFMHMyHDEL.

Cloning of the mouse mannosidase IB catalytic domain with concomitant
 20 addition of the coding sequence for a C-terminal HDEL-tag was done by PCR on a mouse cDNA library (mRNA isolated from the L929 cell line induced with cycloheximide and mouse Tumor Necrosis Factor. Average insert length of the cDNA library was 2000 bp). The PCR oligonucleotide primers used were:
 5'AACTCGAGATGGACTCTTCAAAACACAAACGC3' (SEQ ID NO: 6) and
 25 5'TTGCGGCCGCTTACAACTCGTCGTGTCGGACAGCAGGATTACCTGA3' (SEQ ID NO: 7). The product contained a 5' *Xho* I site and the coding sequence for C-terminal HDEL-site, followed by a stop codon and a *Not* I site at the 3' end. The product was cloned in pGAPZ α A via the *Xho* I/*Not* I sites in the PCR product and the vector, resulting in an in frame fusion of the mouse mannosidase catalytic domain

with the *S. cerevisiae* α -mating factor signal sequence. The sequence of the entire open reading frame generated is set forth in SEQ ID NO: 15.

1.2 Yeast Transformation and Genomic Integration

5

Table 2

Parental strain	DNA transformed
GS115 (his4)	pGAPZMFManHDEL
	pPIC9MFManHDEL
	pPIC9mManHDEL
	pPIC9mMycManHDEL
	pGAPZmManHDEL
	pGAPZmMycManHDEL
GS115 (his4 complemented by pPIC9InfluenzaHA)	pGAPZMFManHDEL
	pGAPZmManHDEL
	pGAPZmMycManHDEL
PPY120H (his4 complemented by pPIC9sOCH1)	pGAPZMFManMycHDEL
	pPICZBMFManMycHDEL
yGC4 (his4 arg1 ade2 ura3 complemented by pBLURA5'PpOCH1)	pPIC9InfluenzaNeuraminidase
	pGAPZMFManHDEL
	pPIC9Glucoseoxidase

All transformations to *Pichia pastoris* were performed with electroporation according to the directions of Invitrogen. Transformants of vectors carrying the Zeocin resistance gene were selected on YPD containing 100 μ g/ml Zeocine

(Invitrogen, Baarn, the Netherlands) and 1M sorbitol. Selection of transformants of pPIC9 derivatives was done on minimal medium lacking histidine and containing 1M sorbitol. Genomic integration of the expression cassettes was verified using PCR on genomic DNA purified from the *Pichia* strains using the Yeast Miniprep method (Nucleon). In all cases concerning the *Trichoderma reesei* gene fusions, the primers used were the sense primer 5'-CCATTGAGGACGCATGCCGCGCC-3' (SEQ ID NO: 8), which annealed to the 3' half of the mannosidase ORF, and the antisense primer 3' AOXI 5'-GCAAATGGCATTCTGACATCCT-3' (SEQ ID NO: 9), which annealed to the AOXI transcription terminator that was present in all our expression constructs.

For the control of genomic integration of the mouse mannosidase transgenes, PCR was done using the sense primer 5'GAP 5'GTCCCTATTTCATCAATTGAA3' (SEQ ID NO: 10, annealing to the GAP promoter or 5'AOXI 5'GACTGGTTCCAATTGACAAGC3' (SEQ ID NO:11), annealing to AOXI promoter), and the antisense primer 3'AOXI (above). For the expression constructs containing a Myc tagged *Trichoderma reesei* α -1,2-mannosidase expression unit, further evidence for genomic integration was obtained using Southern Blotting with the entire MFManMycHDEL ORF (³²P labelled using HighPrime, Boehringer Mannheim) as a probe.

1.3 Expression of α -1,2-mannosidase

Expression of an α -1,2-Mannosidase in GS115 strains expressing influenza virus haemagglutinin was verified by qualitative Northern blot. Expression of an α -1,2-Mannosidase in PPY120H strains was verified by anti-Myc Western blot.

Qualitative Northern Blot -- Total RNA was purified from *Pichia* strains and the yield was determined spectrophotometrically. Northern blotting was performed according to standard procedures and an estimate of the quantity of RNA loaded was made using methylene blue staining of the blot, visualizing the rRNA

bands. The blot was probed with a *Clal/NarI* fragment of the mannosidase, labelled with ^{32}P using HighPrime (Boehringer Mannheim).

SDS-PAGE and Western Blotting -- Total yeast cell lysates were prepared by washing the cells twice with PBS, followed by boiling in 1 volume of 2x concentrated Laemmli loading buffer for 5 min. The lysate was spun briefly in a microcentrifuge prior to gel loading and only the supernatant was loaded. For the analysis of proteins secreted into the growth media, the proteins were precipitated from 200 μl of these media using desoxycholate/trichloroacetic acid according to standard procedures. The pellet was redissolved in 2x concentrated Laemmli loading buffer and the solutions were pH-corrected using Tris. SDS-PAGE was performed and followed by semidry electroblotting to nitrocellulose membranes. For Western Blotting, the 9E10 anti-Myc and the anti-HA mouse monoclonals (Boehringer Mannheim) were used at a concentration of 1 $\mu\text{g/ml}$, and the rabbit anti-PDI antiserum (Stressgen) was used at a dilution of 1/500. The secondary antibodies were goat anti-mouse IgG conjugated to alkaline phosphatase for the monoclonals and goat anti-rabbit IgG conjugated to peroxidase for the polyclonal (secondary antibodies from Sigma). Detection was performed using the NBT/BCIP system for alkaline phosphatase and the Renaissance substrate (NENBiosciences) for peroxidase. Imaging of the latter blot result was done on a Lumilager imaging device (Boehringer Mannheim).

The results shown in **Figure 4** indicated that the great majority of the HDEL-tagged protein was retained intracellularly, both when expressed from the strong constitutive GAP promoter and when expressed from the strong inducible AOXI promoter.

1.4 Localization of α -1,2-Mannosidase

Isopycnic sucrose density gradient centrifugation -- To determine the localization of the HDEL-tagged mannosidase, subcellular fractionation was carried

out using cells expressing the mannosidase-Myc-HDEL from the strong constitutive GAP promoter.

Briefly, 0.5 g of wet weight yeast cells were lysed using 4 x 1 min vortexing with 4.5 g glass beads in 1 ml lysis-buffer (50 mM Tris-HCL pH 7.5 containing 0.6 M sorbitol, 10 mM β -mercaptoethanol and 5 mM $MgCl_2$). Between vortexing periods, the mixture was placed on ice for 5 min. The supernatant was collected and the glass beads were washed once with lysis-buffer, and the supernatant of this washing step was added to the first supernatant. This lysate was subjected to a differential centrifugation procedure. The P10000 pellet was solubilized in 0.5 ml of a 60% sucrose solution in lysis buffer. This solution was placed at the bottom of an Ultraclear ultracentrifuge tube (Beckman) of 14 x 89 mm. Subsequently, 1.5 ml each of sucrose solutions of 55, 50, 45, 42.5, 40, and 37.5% were carefully layered over each other. The tube was filled to the edge with 35% sucrose. Isopycnic sucrose gradient centrifugation was performed for 14 h at 180,000g in a Beckman SW 41 rotor in a Beckman Model L8-70 preparative ultracentrifuge. After completion, 1ml fractions were collected from the top and partially dialysed from excess sucrose, evaporated to dryness in a vacuum centrifuge. After redissolving the pellet in Laemmli buffer, the samples were subjected to SDS-PAGE in triplicate and the Western blots were treated with anti-HA, anti-Myc or anti-PDI ("PDI" for Protein Disulfide Isomerase), respectively.

The results illustrated almost exact cosedimentation of the MFManMycHDEL protein with the Protein Disulfide Isomerase marker protein (which is also targeted with a HDEL signal sequence) (Figure 5). In the same assay, the HA-tagged OCH1 was distributed over the whole gradient, with the highest abundance in fractions having a density lower than that of the fractions containing the mannosidase and the PDI. This result indicated that the mannosidase was targeted to the expected location (the ER-Golgi boundary) by the addition of an HDEL signal. In contrast, the mannosidase without HDEL, expressed from inducible alcohol oxidase I

promoter (which was of comparable strength as the GAP promoter), was secreted at a high level of about 20 mg/l.

Immunofluorescence microscopy -- To confirm the correct targeting of the mannosidase-Myc-HDEL, an immunofluorescence microscopy experiment was
5 performed.

Briefly, yeast cultures were grown to OD₆₀₀ in YPD (for pGAPZMFMManMycHDEL) or in YMP following a YPGlycerol growth phase for pPICZBMFMManMycHDEL. Formaldehyde was added to the yeast cultures to a final concentration of 4% and incubated for 10 min at room temperature. Cells were
10 pelleted and resuspended in 50mM potassium phosphate buffer pH 6.5 containing 1mM MgCl₂ and 4% formaldehyde and incubated for 2h at room temperature. After pelleting, the cells were resuspended to an OD₆₀₀=10 in 100 mM potassium phosphate buffer pH 7.5 containing 1mM MgCl₂ and EDTA-free CompleteTM protease inhibitor cocktail (Boehringer Mannheim). To 100 µl of cell suspension, 0.6 µl of β-mercapto-
15 ethanol and 20µl of 20,000 U/ml Zymolyase 100T (ICN) were added, followed by a 25 minute incubation with gentle shaking. The cells were washed twice in the incubation buffer and added to poly-lysine coated cover slips (these are prepared using adhesive rings normally in use for reinforcing perforations in paper). Excess liquid was blotted with a cotton swab and the cells were allowed to dry at 20°C. All
20 blocking, antibody incubation and washing steps are performed in PBS containing 0.05% bovine serum albumin. Primary antibodies are used at 2µg/µl and secondary antibodies conjugated to fluorphores (Molecular probes) were used at 5µg/µl. The nucleus was stained with the nucleic acid stain HOECHST 33258. After fixation and cell wall permeabilization, the integrity of the yeast cell morphology was checked in
25 phase contrast microscopy and after immunostaining, the slides were examined under a Zeiss Axiophot fluoroescence microscope equipped with a Kodak digital camera. Images were processed using Macprobe 4.0 software and prepared with Corel Photopaint 9.0.

The Golgi marker protein OCH1-HA gave the typical Golgi staining pattern described in the literature (speckle-like staining). Staining with the 9E10 monoclonal anti-Myc antibody, recognizing mannosidase-Myc-HDEL, gave a perinuclear staining pattern with some disparate staining in the cytoplasm, highly
5 indicative for an ER targeting (Figure 4).

Based on the foregoing experiments, it is concluded that the *Trichoderma reesei* mannosidase-Myc-HDEL was targeted to the ER-Golgi boundary.

Example 2

Co-expression of Mannosidase-HDEL 10 with Recombinant Glycoproteins

Co-expression of Mannosidase-HDEL with the *Trypanosoma cruzi trans*-Sialidase

The cloning of a *Trypanosoma cruzi trans*-sialidase gene coding for an active *trans*-sialidase member without the C-terminal repeat domain has been
15 described by Laroy et al. (*Protein Expression and Purification* 20: 389, 2000) which is incorporated herein by reference. The sequence of this *Trypanosoma cruzi trans*-sialidase gene is available through NCBI Genbank under the Accession No. AJ276679. For expression in *P. pastoris*, the entire gene was cloned in pHILD2 (Invitrogen, San Diego, CA), creating pHILD2-TS. To allow better secretion, pPIC9-
20 TS was created in which *trans*-sialidase was linked to the prepro secretion signal of the yeast α -mating factor. Plasmids pPIC9-TSE and pCAGGS-prepro-TSE were created where the epitope E-tag was added to the C-terminal of the *trans*-sialidase to allow easy detection and purification. The construction of pHILD2-TS, pPIC9-TSE and pCAGGS-prepro-TSE has been described by Laroy et al. (2000), incorporated
25 herein by reference. The vectors used in the construction were made available through <http://www.belspo.be/bccm/lmbp.htm#main> for pCAGGS (No. LMBP 2453), Invitrogen, San Diego, CA for pHILD2 and pPIC9, and Pharmacia Biotech for pCANTAB-5E.

Plasmid pPIC9-TSE was linearized with SstI and was transformed into *P. pastoris* GS115 (*his4*) strain by electroporation according to the manufacturer's instructions (Invitrogen). One of the transformants was further transformed with plasmid pGAPZMFManHDEL, establishing a strain co-expressing Mannosidase-HDEL and the *Trypanosoma cruzi* *trans*-sialidase.

Fermentation and protein purification was according to the procedures described by Laroy et al. (2000).

Purified *trans*-sialidase was subject to carbohydrate analysis according to Callewaert et al., *Glycobiology* 11, 4, 275-281, 2001. Briefly, the glycoproteins were bound to the PVDF membrane in the wells of a 96-well plate, reduced, alkylated and submitted to peptide-N-glycosidase F deglycosylation. The glycans were derivatised with 8-amino-1,3,6-pyrenetrisulfonic acid by reductive amination. Subsequently, the excess free label was removed using Sephadex G10-packed spin columns and the glycans were analysed by electrophoresis on a 36 cm sequencing gel on an ABI 377A DNA-sequencer and detected using the built-in argon laser. Digests with 3 mU/ml purified *T. reesei* α -1,2-mannosidase (described by Maras et al., *J. Biotechnol.* 77, 255-63, 2000) were also performed in 20 mM sodium acetate pH=5.0. The glycans derived from 1 μ g of the purified recombinant glycoproteins were used as the substrate. 1U of the α -1,2-mannosidase is defined as the amount of enzyme that releases 1 μ mol of mannose from baker's yeast mannan per minute at 37°C and pH=5.0.

As can be seen in Figure 6, panel B, the major N-glycan on *trans*-sialidase was Man₈GlcNAc₂ (Compare with panel F, representing an analysis of the N-glycans of bovine RNaseB. The one but last peak in this profile is Man₈GlcNAc₂, the first peak is Man₅GlcNAc₂). *In vitro*, this glycan was digestible to Man₅GlcNAc₂ with α -1,2-mannosidase (Figure 6, panel C). In the N-glycan profile of the *trans*-sialidase co-expressed with mannosidase-HDEL, the major peak corresponded to Man₅GlcNAc₂ (Figure 6, panel D).

Co-expression of Mannosidase-HDEL with the Influenza A virus haemagglutinin

The Influenza A virus haemagglutinin was known to be glycosylated in *Pichia pastoris* with high-mannose N-glycans containing 9-12 mannose residues (Saelens et al. *Eur. J. Biochem.* 260: 166-175, 1999). The effect of a co-expressed
5 mannosidase on the N-glycans of the haemagglutinin was assessed in an N-glycan profiling method described below. In addition, to compare the efficiency of the *Trichoderma* enzyme (having a temperature optimum of 60°C) with a mammalian mannosidase having a temperature optimum of 37°C, the catalytic domain of the mouse mannosidase IB from a mouse cDNA-library was cloned and tagged with a
10 HDEL signal by PCR amplification. This ORF was cloned after the prepro-signal sequence of the *S. cerevisiae* α -mating factor under the control of the GAP promoter. Expression of the mannosidase-HDEL transgenes on the mRNA level was confirmed by qualitative Northern blotting.

The haemagglutinin was expressed and purified from a non-mannosidase
15 expressing control strain and from a strains co-expressing the *Trichoderma reesei* mannosidase-HDEL or the mouse mannosidase IB-HDEL according to the procedure described by Kulakosky et al. *Glycobiology* 8: 741-745 (1998). The purified haemagglutinin was subjected to PNGase F digestion as described by Saelens et al. *Eur. J. Biochem.* 260: 166-175, 1999. The proteins and glycans were precipitated with 3
20 volumes of ice-cold acetone and the glycans were extracted from the pellet with 60% methanol. Following vacuum evaporation, the glycans were labeled with 8-amino-1,3,6 pyrenetrisulfonic acid by adding 1 μ l of a 1:1 mixture of 20 mM APTS in 1.2M citric acid and 1M NaCNBH_3 in DMSO and incubating for 16h at 37°C at the bottom of a 250 μ l PCR-tube. The reaction was stopped by the addition of 10 μ l deionized
25 water and the mixture was loaded on a 1.2 cm Sephadex G10 bed packed to dryness in a microspin-column by centrifugation in a swinging bucket rotor, which provided for a flat resin surface. After loading, 50 μ l deionised water was carefully added to the resin bed and the spin column was briefly centrifuged for 5 seconds at 750g in a tabletop centrifuge. This elution process was repeated twice and all the eluates were

pooled and evaporated to dryness in a Speedvac vacuum centrifuge (Savant). The labeled glycans were reconstituted in 1.5 μ l gel loading buffer containing 50% formamide and 0.5 μ l Genescan 500TM, labeled with rhodamine (Perkin Elmer Bioscience), serving as an internal reference standard. This mixture was loaded on a
5 DNA-sequencing gel containing 10% of a 19:1 mixture of acrylamide:bisacrylamide (Biorad, Hercules, CA, USA) and made up in the standard DNA-sequencing buffer (89 mM Tris, 89 mM borate, 2.2 mM EDTA). Polymerization of the gel was catalyzed by the addition of 200 μ l 10% ammonium persulfate solution in water and 20 μ l TEMED. The gel was of the standard 36 cm well-to-read length and was run on
10 an Applied Biosystems Model 373A DNA-sequencing apparatus. Prerunning of the gel was done at 1000 V for 15 min. and after loading, the gel was electrophoresed for 8h at 1250 V without heating. This methodology gives a limit of detection of 10 fmol per peak. The data were analysed with Genescan 3.0 software.

As shown in **Figure 7**, the *Trichoderma reesei* α -1,2-mannosidase
15 provided the most complete reduction in the number of α -1,2-mannoses present on the N-glycans. The N-glycan processing by mouse mannosidase IB-HDEL was less efficient than by the *Trichoderma reesei* α -1,2-mannosidase.

Despite the efficient removal of α -1,2-mannoses from the N-glycans of haemagglutinin, no $\text{Man}_5\text{GlcNAc}_2$ was obtained. Even after digestion of the N-
20 glycans with 3 mU of purified *Trichoderma reesei* α -1,2-mannosidase, only $\text{Man}_6\text{GlcNAc}_2$ was obtained as the smallest sugar chain. These results indicated that the remaining residues were possibly α -1,6-linked mannoses, originating from the initiating OCH1 α -1,6-mannosyltransferase enzymatic activities. OCH1 was observed to be localized to very early part of the Golgi apparatus and could act on the N-glycans
25 of haemagglutinin before complete digestion of the $\text{Man}_8\text{GlcNAc}_2$ precursor to $\text{Man}_5\text{GlcNAc}_2$ by the mannosidases-HDEL. Thus, for proteins whose glycans are efficiently modified by the α -1,6-mannosyltransferase, an inactivation of the OCH1

gene coding for the transferase would be desirable in order to obtain proteins with $\text{Man}_5\text{GlcNAc}_2$.

Example 3

Inactivation of the *Pichia* Och1 Gene:

A *Pichia pastoris* sequence was found in the GenBank under Accession
5 No. E12456 and was described in Japanese Patent Application No. 07145005,
incorporated herein by reference. This sequence shows all typical features of an α -
1,6-mannosyltransferase and is most homologous to the *S. cerevisiae* OCH1, thus
referred to herein as the *Pichia pastoris* Och1 gene.

First, the full ORF of the *Pichia pastoris* Och1 gene was PCR cloned in
10 pUC18 to obtain plasmid pUC18pOch1. pUC18pOch1 was cut with HindIII, blunt-
ended with T4 polymerase, then cut with XbaI, releasing a fragment containing the 5'
part of the *Pichia pastoris* Och1 gene. This fragment was ligated into the vector
pBLURA IX (available from the Keck Graduate Institute, Dr. James Cregg,
<http://www.kgi.edu/html/noncore/faculty/cregg/cregg.htm>), which had been cut with
15 *Eco* RI, blunt-ended with T4 polymerase, and then cut with *Nhe* I. This ligation
generated pBLURA5'PpPCH1, as shown in **Figure 8**.

Disruption of this *Pichia* OCH1 gene in the *Pichia* genome was achieved
by single homologous recombination using pBLURA5'PpOCH1, as illustrated in
Figure 9. As a result of the single homologous recombination, the Och1 gene on the
20 *Pichia* chromosome was replaced with two Och1 sequences: one consisted only about
the first one third of the full Och1 ORF, the other had a full Och1 ORF without a Och1
promoter. Single homologous recombination was achieved as follows. Cells of the
Pichia strain yGC4 were transformed by electroporation with pBLURA5'PpOCH1
which had been linearized with the single cutter *Bst* BI. About 500 transformants
25 were obtained on minimal medium containing 1M sorbitol, biotin, arginine, adenine
and histidine and incubation at 27°C. Thirty-two of these transformants were picked
and re-selected under the same conditions. Twelve clones were further analyzed for
correct genomic integration of the cassette by PCR. Seven of the twelve URA
prototrophic clones contained the cassette in the correct locus.

One of the Och1-inactivated clones was also further transformed with pGAPZMFManHDEL to produce "supertransformants". Both the Och1-inactivated clone and three supertransformants also expressing the ManHDEL were evaluated in cell wall glycan analysis as follows. Yeast cells were grown in 10 ml YPD to an

5 OD₆₀₀=2 and mannoproteins were prepared by autoclaving the yeast cells in 20 mM sodium citrate buffer pH7 for 90 min at 120 °C and recovery of the supernatant after centrifugation. Proteins were precipitated from this supernatant with 3 volumes of cold methanol. The protein preparation obtained in this way was used for N-glycan analysis using DSA-FACE as described by Callewaert et al. (2001) *Glycobiology* 11,

10 275-281. As shown in **Figure 10**, there was an increased amount of Man₈GlcNAc₂ glycan in the Och1-inactivated clone as compared to parent strain yGC4, indicative of a reduced activity of the Och1 enzyme. In all three supertransformants which also expressed the HDEL-tagged α -1,2 mannosidase, the production of Man₅GlcNAc₂ was observed. Furthermore, upon digestion of the same glycan mixtures with 3 mU/ml

15 purified recombinant *Trichoderma reesei* α -1,2-mannosidase, more Man₅GlcNAc₂ was formed in the strain transformed with pBLURA5'PpOCH1 than in the parent strain (**Figure 11**, compare panel 2 and 3).

These results confirmed that the lack of a production of Man₅ glycans on recombinantly produced proteins such as haemagglutinin from cells expressing α -1,2-

20 mannosidase were due to the activity of the Och1 protein. These results further indicate that the production of glycoproteins with Man₅ glycans could be facilitated by the inactivation of the Och1 gene.

Example 4**Expression of Glucosidase II in *Pichia pastoris*****4.1 Amplification of the GLSII alpha subunit ORF from *S. cerevisiae*.**

5 Genomic DNA was prepared from the *S. cerevisiae* strain INVS (α , leu2-3, 112 his3 Δ 1, trp1-289, ura3-52), using the Nucleon kit (Amersham). A touch-down PCR reaction was performed using this genomic DNA as template and the LA TaKaRa polymerase (ImTec Diagnostics). The sequence of the PCR primers was based on the known sequence of the *S. cerevisiae* GLSII ORF:

10 Sense primer: 5' CCG CTC GAG ATG GTC CTT TTG AAA TGG CTC 3'
Xho I

(SEQ ID NO:12)

Antisense primer: 5' CCG GGC CCA AAA ATA ACT TCC CAA TCT TCA
Apa I

15 G 3' (SEQ ID NO:13)

4.2 Cloning of the *S. cerevisiae* glucosidase II ORF into *Pichia pastoris* expression vectors.

Construction of the glucosidase II expression vectors -- The PCR fragment
 20 was digested with *Xho* I/*Apa* I and ligated into the pGAPZA vector (Invitrogen), thereby placing the ORF under the transcriptional control of the GAP promoter. Using this strategy, the myc and the His6 tag were placed in frame to the C-terminus of Glucosidase II, creating pGAPZAGLSII. The complete ORF of pGAPZAGLSII was then sequenced to ensure that no mutations were generated in the PCR reaction.
 25 The sequence of the vector pGAPZAGLSII was set forth in SEQ ID NO: 18. The GLSII ORF from the pGAPZAGLSII vector was cloned into vector pPICZA (Invitrogen) to create pPICZAGLSII, thereby placing the ORF under the transcriptional control of the AOXI promoter. The GLSII ORF from the pGAPZAGLSII vector was cloned into vector pAOX2ZA, thereby placing the ORF
 30 under the transcriptional control of the AOX2 promoter. This vector was created by

replacing the multi cloning site of vector pAOX2ZB with the multi cloning site of pPICZA. Vector pAOX2ZB was generated by replacing the AOX1 promotor of pPICZB by the AOX2 promotor region of the AOX2 gene (Martinet et al., Biotechnology Letters 21). The AOX2 promotor region was generated by PCR on

5 *Pichia* genomic DNA with the sense primer 5'GACGAGATCTTTTTTTCAGACCATATGACCGG 3' (SEQ ID NO: 26) and the antisense primer 5'GCGGAATTCTTTTCTCAGTTGATTTGTTTGT 3' (SEQ ID NO: 27). The GLSII ORF from the pGAPZGLSII vector was cloned into vector pYPT1ZA to create pYPT1ZAGLSII, thereby placing the ORF under the

10 transcriptional control of the YPT1 promoter. Vector pYPTZA was created by replacing the AOX1 promoter of pPICZA by the YPT1 promoter present on the plasmid pIB3 (GenBank accession number AF027960)(Sears et al., Yeast 14, pg 783-790, 1998). All constructs contain the phleomycin resistance gene. The resulting final expression vectors (pGAPZAGLSII, pAOX2ZAGLSII, pPICZAGLSII and

15 pYPT1ZAGLSII) are depicted in **Figures 12-15**.

Similar expression vectors were constructed, carrying the Ampicillin resistance marker and the *Pichia* ADE1 selection marker. In principle, the Zeocin resistance expression cassette of the plasmids pAOX2ZAGLSII, pGAPZAGLSII and pYPT1ZAGLSII was replaced by the Ampicillin and *Pichia* ADE1 cassette of the

20 vector pBLADE IX (Cregg, J.M.) to result in the vectors pAOX2ADE1glsII, pGAPADE1glsII and pYPT1ADE1glsII. Vector pPICADE1glsII was obtained by inserting the glucosidase II open reading frame into the multiple cloning site of the vector pBLADE IX (Cregg, J.M.). The resulting final expression vectors (pGAPADE1glsII, pAOX2ADE1glsII, pPICADE1glsII and pYPT1ADE1glsII) are

25 depicted in **Figures 16-20**.

Adding the ER retention tag HDEL to Glucosidase II expression vectors –

The following primers were used to generate an HDEL-containing PCR fragment:

Primer 1: 5'GCG GGT CGA C/CA C/GA C/GA A/CT G/TG A/GT TTT AGC CTT
 Sal I H D E L stop

AGA CAT GAC 3' (SEQ ID NO:28)
Primer 2: 5'CAG GAG CAAA GCT CGT ACG AG 3' (SEQ ID NO:29)
*Spl*I

5 PCR was performed on pGAPZAGLSII with Taq pol., at 60°C. The PCR
fragment of 225 bp was cut with *Sal* I/*Spl* I and ligated into the *Sal* I/*Spl* I opened
pGAPZAGLSII vector, creating plasmid pGAPZAglslIIHDEL. The sequence of
plasmid pGAPZAglslIIHDEL is set forth in SEQ ID NO: 24. The construction
strategy and the resulting final expression vectors (pGAPZAglslIIHDEL and
10 pGAPADElglslIIHDEL) are depicted in **Figures 20-21**.

4.3 Transformation of a *Pichia pastoris* strain.

Transformation was performed using the conventional electroporation techniques, as described by Invitrogen. Cells of the *Pichia pastoris* strain PPY12-OH
15 were transformed with pGAPZGLSII which had been cut with the single cutter *Avr* II. Transformants were selected based on their resistance to zeocin.

Genomic analysis of the transformants -- Genomic DNA was prepared from some zeocin resistant *Pichia* transformants. A PCR reaction was performed on the genomic DNA in order to determine whether or not the glucosidase II gene was integrated into the yeast genome. PCR was performed using Taq DNA polymerase (Boehinger) (2.5 mM MgCl₂, 55°C for annealing). The primers were the same as the ones we used for the amplification of the ORF on *S. cerevisiae* genomic DNA. pGAPZAGLSII transformants were confirmed by the presence of a specific PCR product indicative of the glucosidase II ORF.

4.4 Expression and secretion of the *S. cerevisiae* glucosidase II alpha subunit in *Pichia pastoris*

Analysis at the transcriptional level -- RNA was prepared from the transformants which scored positive after the genomic analysis. RNA was prepared using acid phenol. From each sample, 15 µg of RNA was loaded on a formaldehyde

agarose gel. After electrophoresis the RNA was blotted on a Hybond N membrane. The membrane was hybridizing using a radioactive probe, which consists of a 344 bp glucosidase II specific fragment, corresponding to the 3' region of the glucosidase II ORF. No signals were detected with non-transformed control strains, whereas clear
5 signals were observed with transformants.

Analysis at the protein level using a double membrane assay -- A nitrocellulose membrane was placed on a buffered dextrose medium (BMDY). On top of that nitrocellulose membrane, a cellulose acetate membrane was placed. *Pichia* transformants of pGAPZAGLSII were streaked on the cellulose acetate and grown for
10 a few days. The yeast cells remained on the cellulose acetate, while the secreted proteins crossed this membrane. As such the secreted protein was captured onto the nitrocellulose membrane. After a few days the cellulose acetate, containing the yeast colonies, was removed. The nitrocellulose membrane was analyzed for the presence of glucosidase II using anti-myc antibody. Most of the transformants gave a clear
15 signal as compared to a faint, hardly visible signal with the WT, non-transformed strain.

Extracellular expression -- PPY12-OH transformants of the construct pGAPZAGLSII(myhis6) (strains 12, 14 and 18) and transformants of the construct pGAPZAGLSII(myc)HDEL (strains H1, H2 and H3) were grown for 2 days on 2x10
20 ml BMDY medium. These 6 transformants earlier scored positive both on the genomic level (PCR on gDNA) and on the RNA level (Northern blot). The culture medium was collected by centrifugation and concentrated with Vivaspin columns to about 1 ml. Proteins from this concentrate were precipitated with TCA, resuspended in Laemmli buffer and loaded for SDS-PAGE analysis. Proteins were blotted to
25 nitrocellulose membrane. The blot was incubated overnight with anti-myc Ab. The secondary Ab was linked to peroxidase. Using the Renaissance luminiscence detection kit (NEN) and a light sensitive film (Kodak), a strong band at about 110 kDa was observed for the transformants 12, 14 and 18, indicating that GLSII was expressed and secreted from these transformants. No signal was obtained for the transformants

H1-3, which indicate that the HDEL tag, which was added C-terminally to the GLSII ORF, resulted in an ER localization of the protein, preventing GLSII to be secreted into the growth medium.

Intracellular expression – The 6 transformants and the WT strain were grown for 2 days in 500 ml BMDY. The cells were collected by centrifugation, washed, resuspended into a minimal volume (50 mM Tris.HCl pH 7.5, 5% glycerol) and broken using glass beads. The cell debris was removed through several centrifugation steps (low speed centrifugation (2000-3000g)). Membranes were obtained from the supernatant through ultracentrifugation. The pellets were resuspended in Laemmli buffer and loaded for SDS-PAGE analysis. The proteins were blotted on a nitrocellulose membrane. The intracellular GLSII expression was checked using anti-myc Ab and peroxidase conjugated secondary Ab. Following the luminescence detection, a band at about 110 kDa was observed with the GLSIIHDEL transformants (H1 and H3, faint signal for H2), but not with the WT and GLSII expression strains. These results clearly indicate the intracellular presence of the recombinant GLSII when expressed with a C-terminal HDEL tag. No GLSII was detected intracellularly when this tag was not present.

4.5 Purification and activity assays of the recombinant glucosidase II alpha

submit

A GLSII assay was set up as follows and was tested using a commercially available yeast alpha-glucosidase (Sigma) as a positive control.

Composition: 70 μ l 80 mM phosphate-citrate buffer pH 6.8, 7 μ l 250 mM mannose, 3.5 μ l 250 mM 2-deoxy-D-glucose, 0.8 μ l 4-MeUmbelliferyl-alpha-D-glucopyranoside (1 μ M). Three assays were performed: one with 1 unit commercial enzyme, one without the enzyme and one with the enzyme but without the substrate. The assay mixture was incubated overnight at 30°C. When illuminated with UV, only the reaction mixture with both the enzyme and the substrate showed fluorescence

(Figure 22). This indicates that the assay was very specific in detecting the activity of the alpha-glucosidase.

- WT PPY12-OH, strain 18 and strain H3 were grown during 2 days in 2x10 ml growth medium. Cells were spun down and medium was adjusted to 300 mM NaCl and 10 mM imidazol and concentrated with Vivaspin columns to 0.5-1ml. Medium was loaded onto a Ni-NTA spin column (Qiagen) and the purification was performed according to the manufactures recommendations. Protein was eluted from the column in 2x100 µl elution buffer (50 mM NaH₂PO₄, 300 mM NaCl, 250 mM imidazol pH 8.0). From each eluate, 20 µl was assayed for its glucosidase II activity.
- 0.33 units of the commercial enzyme diluted in 20 µl of the elution buffer was used as a positive control. Fluorescence was observed with the positive control and the elute of strain 18, the strain which secreted the enzyme into the growth medium. These results indicate that the recombinant *S. cerevisiae* GLSII alpha subunit, secreted by *Pichia pastoris*, was a functionally active enzyme. The activity was not seen in the WT (untransformed) strain, nor in strain H3 as the GLSII was expressed intracellularly (Figure 23). These results also indicate that the beta subunit is not necessary for the functionality of the alpha part of the protein.

Example 5**Creating *Pichia* Strains Expressing
both Glucosidase II and Mannosidase**

5 Strain GS115 was transformed with pGAPZGLSII and pGAPZglsIIHDEL.
Transformants were selected on YPDSzeo.

Strain yGC4 was transformed with the following constructs, respectively:

- (1) pGAPADEglsII and pGAPADEglsIIHDEL, selection on synthetic
sorbitol medium without adenine;
10 (2) pGAPZMFManHDEL: selection on YPDSzeo; and
 (3) pGAPZMFManHDEL/pGAPADEglsIIHDEL: selection on synthetic
sorbitol medium without adenine and with zeocin.

Strain yGC4 with OCH1 knock-in and expressing MFmannosidaseHDEL
was transformed with pGAPADEglsII and pGAPADEglsIIHDEL. Selection of
15 transformants was done on synthetic sorbitol medium without adenine and uracil.

For all transformations, colonies were obtained. Transformants with the
expression vector(s) integrated into the genome, determined by PCR, were obtained.
Expression of GLSII from some of these transformants was observed.

20

SEQUENCE LISTING

- 5 **SEQ ID NO: 1**
HDEL (peptide)
- SEQ ID NO: 2**
5'-GACTGGTTCCAATTGACAAGC-3'
- SEQ ID NO: 3**
5'-AGTCTAGATTACAACCTCGTCGTGAGCAAGGTGGCCGCCCCG TCG-3'
- 10 **SEQ ID NO: 4**
CCATTGAGGACGCATGCCGCGCC
- SEQ ID NO: 5**
GTATCTAGATTACAACCTCGTCGTGCAGATCCTCTTCTGAGATGAGTTTTTGT
15 TCAGCAAGGTGGCCGCCCCGTCGTGATGATGAA
- SEQ ID NO: 6**
AACTCGAGATGGACTCTTCAAAACACAAACGC
- 20 **SEQ ID NO: 7**
TTGCGGCCGCTTACAACCTCGTCGTGTCGGACAGCAGGATTACCTGA
- SEQ ID NO: 8**
CCATTGAGGACGCATGCCGCGCC
- 25 **SEQ ID NO: 9**
GCAAATGGCATTCTGACATCCT
- SEQ ID NO: 10**
30 GTCCCTATTTCAATCAATTGAA
- SEQ ID NO: 11**
GACTGGTTCCAATTGACAAGC
- 35 **SEQ ID NO: 12**
CCG CTC GAG ATG GTC CTT TTG AAA TGG CTC
- SEQ ID NO: 13**
40 CCG GGC CCA AAA ATA ACT TCC CAA TCT TCAG

gcacaaataacgggtattgttataaatactactattgccagcattgctgctaagaagaagggtatctctcgagatggact
cttcaaacacaaacgcttggatctgggcttagaagatgtgttaattcctcacgtagatgccggcaaggagctaaaaacccc
ggcgtcttcctgatccatggacccgacgaacacagacacagggaagaagaagagcgtctgagaaataagattagagctg
accatgagaaagccctggaagaagcaaaagaaaaaattaagaaagtcaagagaggaaatccgtgcagaaattcagacaga
5 gaaaaacaaagtagcccaagcaatgaagacaaaagagaccagggtactgccgcctgtccctgtcccacaacgtgtaggg
gtcagtggtggggatccagaagacatggagatcaagaagaaaagagacaaaattaaagagatgatgaacatgcctggg
ataaltacagaacatacggatggggacataatgaactaaggccatttgaaggaaggccattccactaacatattcggag
ctcacagatgggtgccaccatagtggtgcttggataccctttatatcatggggcttcatgatgaattcatggatgggcaag
atggattgaagaaaccttgatttcagtggaattcagaagtgtctgtcttgaagtaacattcgtttattggagggtcctcgc
10 ctgcatattacctgtcaggagaggaatattcaagactaaagcagtgagtggtgctgagaaactcctcctgccttaacacac
ctactgggattcctggcaatgggtgaacctgaaaagtggagtaggtcgaactggggctgggcgtctgcaggcagcagc
atcctggctgagttcggcaccctgcacatggagttgtgcacctcagctacttgaccgggtgacttgacttactataataaggtc
atgcacattcggaaactactgcagaaaatggaacgcccaatggctttatccaaattatttaaaccgaacaggcgctg
gggtcagtatcacacatcagttggtggtctgggagatagttttatgaatacttactgaaagcatggctgacgtcagataaac
15 agaccacgaggcaagaaggatgtatgacgatgctgtgaggctatagaaaaacatcttattaagaagtcccgaggaggtct
gggttttattggagaatggaagaatggacacttggaaagggaagatggggcacttggcctgcttggctgggggaatgcttgc
cttggagcagatggtccagaaaggataaagctggccactacttagaactaggggcagaaattgcacgaacatgtcatgag
tcatatgacagaactgcattgaaactagggtccggagtcattcaagttgatggtgcagtggagccgtggctgtgcggcagg
ctgaaaagtattacatccttcgtccagaagtaattgaacctattggtatctatggcgatttaccacgaccaagatacaggc
20 agtggggctgggaagcagcactggctattgagaagtcgtccgggtcagcgggtgggtttctggtgtcaaggatgtatag
ccccgacctgtgcatgacgacgtgcagcagagctttctctgtgaaacattaaaatactgtacctgtgttctctggcga
tgaccttctaccttagaccactgggtgttaacacagaggcgccacctctgcgggtgttgcgcttagccaacagcactcttc
aggtaatcctgctgtccgacacgacgagttgtaa

25 **SEQ ID NO: 16** pAOX2ZAGLSII

catggccaagtggaccagtgccgttcgggtgtcaccgcgcgcagctgcgggagcgggtcaggttctggaccgaccgg
ctcgggttctccgggacttctggaggacgacttcgccgggtgtggtccgggacgacgtgacctgttcatcagcgcggtc
caggaccaggtggtgccgacaacacctggcctgggtgtgggtgcgcggcctggacgagctgtacccgagtggtcgcg
gaggtcgtgtccacgaactccgggacgcctccgggcccggccatgaccgagatcggcgagcagccgtggggggcggga

gtctgcccctgcgcgacccggccggcaactgcgtgcacttcgtggccgaggagcaggactgacacgtccgacggcgccc
 cacgggtcccaggcctcgagatccgtcccccttttcttctgcatatcatgtaattagttatgtcacgttacatcacgccc
 cccccacatccgclctaaccgaaaaggaaggagtagacaacctgaagtctaggtccctattttttatagttatgttagt
 attaagaacgtatttataatttcaatttttctttttctgtacagacgcgtgtacgcatgtaacattatactgaaaaccttgctga
 5 gaagggtttgggacgctcgaaggctttaatttgcaagctggagaccaacatgtgagcaaaaggccagcaaaaggccagga
 accgtaaaaaggccgcttgctggcgttttccataggctccgccccctgacgagcatcacaanaatcgacgctcaagtca
 gaggtggcgaaacccgacagactataaagataccaggcggttccccctggaaagctccctcgtgcgctcctctgtccgac
 cctgcccgttacgggatacctgtccgcttttctcccttcgggaagcgtggcgctttctcaatgctcacgctgtaggtatctcagt
 tcggtgtaggtcgtcgtccaagctgggctgtgtgcacgaacccccgttcagcccagccgctgcgcttatccggtaact
 10 atcgtcttgagccaacccggtaagacacgactatcggcactggcagcagccactggtaacaggattagcagagcgaggt
 atgtaggcggtgtacagagttctgaagtgggtggcctaactacggctacactagaaggacagttattggatctgcgctctg
 ctgaagccagttacctcgaaaaagagttggtagctcttgatccggcaaaacaaaccaccgctggtagcggtggtttttgtt
 tgcaagcagcagattacgcgcagaaaaaaggatctcaagaagatccttgatctttctacggggctgacgctcagtgga
 acgaaaaactcacgtaagggttttggcatgagatcagatctttttcagaccatatgaccgggtccatctctacggggggat
 15 tatctatgctttgacctctatcttgattctttatgattcaaatcacctttacgttatttacttactggttatttacttagccctttctg
 aaaaacatttactaaaaatcatacatcggcactctcaaacacgacagattgtgatcaagaagcagagacaatcaccactaag
 gtgtcacatttgagccagtaggctcctaataagaggttcgatacttatttgataatacgacatattgtcttacctctgaatgtgtca
 atactctctcgttctcgtctcgtcagctaaaaatataacacttcgagtaagatacggccaattgaaggctacgagataccaga
 ctatcactagtagaactttgacatctgctaaagcagatcaaatatccattatccagaatcaattaccttcccttagctgtcgaag
 20 gcatgaaaaagctacatgaaaatccccatcctgaagtttgcagcttaaggactccatttctaaaattcaagcagtcctct
 caactaaatttttccattcctctgcaccagccctctcatcaaccgtccagccttctcaaaagtccaatgtaagtagcctgca
 aattcaggttacaacccctcaattttccatccaaggcgatccttacaaggttaataatcgaacagcagagactaagcgagtc
 catcaccaccaacgatgggtgaaaaacttaagcatagattgatggagggtgtatggcacttggcggtgcatlagagttga
 aactatggggaatacatcacatccggaactgatccgactccgagatcatatgcaaagcacgtgatgtacccgtaaaactgc
 25 tcggattatcgttgcaattcatcgtcttaaacagtacaagaaactttatcattgggtcattggactctgataggggcacattcc
 ccaatgatttttgggaaagaaagccgtgaaggagacagttgaagcgaagagacaagacaacgaacagcaaaagtacag
 ctgtcagctacctagtgacaggttgggagtttccaattgggtgtttgaatttttaccatgttgagttgctctctccttgca
 aacaatgcaagttgataagacatcaccttccaagataggctattttgtgcataaattttgtctcgagtgaaaacccctttat
 gtgaacagattacagaagcgtcctaccctcaccgggtgagatggggagaaaattaagcgaatgaggagacgattattggtat

aaaagaagcaaccaaatacccttattgtccttttctgatcagcatcaaagaataattgtctfaaacgggcttttaactacattgttc
 ttacacattgcaaacctcttcttcttattcggatcaactgtattgactacattgatcttttaacgaagttfacgactfactaaatcc
 ccacaaacaaatcaactgagaaaagaattcacgtggcccgccggtctcggatcggtacctcgagatggccttttga
 aatggctcgtatgccaattggcttcttaccgctttttcgcatgcgtttaccgactatctattaagaagtgtgcgaatctgggt
 5 tttgccatagaaacagggtttatgcagaaaatttgccaaatctcatcactgctattacaaagtgagcgagcttattgcac
 acgatcctttagagaatgtgcttcatgctaccataatfaaaactataccaagattggaggcgcatgatatagccgttcagtcc
 attctctctctttttacaggatcactcagtaagggtcactataaatgagaaagagagaatccaaccaacagcagcggttg
 ttgatctcttcacaacgggtcaatgagacctggaagtacgcattcgacaagaaattcaagaggaggcgacaggaccagta
 ttccacaattccacttcttaagcaaaaacaaactgtgaactcatttggctgaaaaatatcttcattttgtcactttcaaacccac
 10 tgcagacacatttcatcttcgaaacgggtgatgtatccgtagaatactttgtgaacctttcaattgaaagtttactggcaaatg
 cgctgaaacttattgtaaacgagcaaaatttctgaacattgaacatcatagaactaagcaggaaaacttcgcacacgtgctg
 ccagaagaacaacttcaacatgtttaaggacaatttctgtatcaagcatgactctatgcccttggggcctgaatcggttg
 cgctagatttcttctcatgggttctactaatgtctacgggtatccggaacatcgacgtcgctaaggctgatggacacttcagg
 tggaaaggaacctacaggcttttcaacgttgatgtcttfgatcaaacatcggtaccagccaaccaatgtacgggttcgatcc
 15 catlcatgttttcatcttcgtccacatctatctttgggtcaatgcagctgacacttgggtagacataaagtatgacaccagtaaa
 aataaaacgatgactcattggatctccgaaaatgggtgcatagatgtatcatgtccctggggccagatattccaactatcattg
 acaatttaccgatttgactggtagacctttttaccgccatttctctatagggtaccatcaatgtagatggaattataatgatg
 agatggacgttctcacagtggtactctcagatggatgctcatatgattccttacgattttatttgggtgacttggagtatacgaac
 gacaaaaaataatttacttggaaagcagcactccttcccaatccaaaaaggctgttatccaaattaaaaaagggtgggtagaat
 20 cttgtcgtactaatcgatcctcatttaaagaaagattatgaaatcagtgacagggttaattaatgaaatgtagcagtcaaggatc
 acaatggaaatgactatgtaggtcattgctggccaggtaattctatatggattgataccataagcaaatatggccaaaagattt
 ggaagtccttttgcgaacgggttatggatctgccggtgatttaactaatitattcatttggatgatatgaacgagccttcgattt
 cgatggccagagaccacagctccaaaagatttgattcacgacaattacattgaggaaagatccgtccataacatatatggtc
 tatcagtgcatgaagctacttacgacgcaataaaatcgatttaccatccgataagcgtccttcttctaacaagggttttt
 25 ttgccggtctcaacgtactgctgccacatggactggtgacaatgtggccaattgggattacttaaaagatttccattcctatggt
 tctgtcaacaacattgctggtatgccatttataggagccgacatagctggcttctgtgaggatcctacacctgaattgattgca
 cgttggtagcaagcgggttatgtaccatttttagagcacacgcccataatagaccaagagaagagaaccatacttatt
 caatgaacctttgaagtcgatagtacgtgatatttcaattgagatatttctgctacctacctatataccatgtttcataaatca
 agtgtcactggatttccgataatgaatccaatgtttatgaacacctgaatttgcgaattgtatcatatcgataaccaatttact

ggagtaattcaggtctattagtcaaacctgtcacggagcctggtaacagaaacggaaatggtttccaccgggtatattct
 atgaattcgcactctttacactctttataaacaatgggtactgattgatagaaagaatatttctgcaccattggataaaattccatt
 atttattgaagcggtcacaltatcactatgaaagataagtatagaagatcttcaatgtaatgaaaaacgatccatatgtaatag
 ttatagcccctgataccgaggagcagcgttggagatcttattgtgatgatggagaaacttttggctaccaagagggtgag
 5 tacgtagaaactcagttcattttcgaacaataccttaaaaaatgttcgaagtcataatcccgagaatttgacaggcattcacc
 acaatactttgaggaataccaatattgaaaaatcattatcgcaagaataattacaacacaacataacgttgaaagacagta
 ttaaagtcaaaaaaatggcgaagaagttcattgccgactagatcgatgagaatgataataagatcaccattcttaacc
 tatcgcttgacataactgaagattgggaagtatttttggcccgaacaaaaactcatctcagaagaggatctgaatagccc
 gtcgaccatcatcatcatcatcattgagtttagccttagacatgactgttcctcagttcaagttgggcacttacgagaagaccg
 10 gtcttgctagattctaatacagaggatgtcagaatgccatttgcctgagagatgcaggcttcattttgatactttttattgtaac
 ctalatagtataggattttttgtcattttgtttctctgtagcagcttgcctgatcagcctatctcgcagctgatgaatatcttgt
 ggtagggggttgggaaatcattcgagtttgatgttttcttggtatttccactcctcttcagagtacagaagatlaagtgagac
 ctctgttgtgcggatccccacacaccatagctcaaaatgtttactcctttttactcttccagattttctcggactccgcga
 tcgccgtaccacttcaaacacccaagcacagcatactaaattttcctcttcttctctaggggtgcgttaattacccgtacta
 15 aagggttggaaaagaaaaagagaccgcctcgttcttttctctcgtcgaaaaaggcaataaaattttatcacgtttctttctt
 gaaatttttttttagtttttcttcttcagtgacctcattgatatttaagtaataaacggcttcaatttctcaagtttcagttcatt
 ttctgttctattacaacttttttacttctgttcattagaagaaaagcatagcaatctaataaggcggtgttgacaattaatcat
 cggcatagtalatcggcatagataatacgaagaaggtaggaactaac

20 **SEQ ID NO: 17 pAOX2ADE1glsII**

tcgaccggctgcattaatgaatcgccaacgcgcggggagaggcggttgcgtattgggcgctcttccgcttctcgtcac
 tgactcgtcgtcgtcgtcgttgcggctgcggcagcgggtatcagctcactcaaggcggtaatacgggtatccacagaatca
 ggggataacgcaggaaagaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccggttgcgtggc
 gttttccataggctccgccccctgacgagcatcacaaaaatcgacgctcaagtcagagggtggcgaacccgacaggact
 25 ataaagataccaggcggtttccccctggaagctccctcgtcgcgtctcctgttccgacctgccgcttaccggatacctgtccg
 cctttctccctcgggaagcgtggcgctttctcalagctcacgctgtaggtatctcagttcgggtgtaggtcgttcgctcaagct
 gggctgtgtgcacgaacccccgttcagcccagccgtcgccttatccggtaactatcgtcttgagtcacccggtaaga
 cagacttatcgtccactggcagcagccactggtacaggattagcagagcgaggatgtaggcgggtgtacagagttcttg
 aagtgggtgcctaactacggctacactagaaggacagatttggatctcgcgtctcgtgaagccagttaccttcggaaaaag

agttggtagctcttgatccggcaacaaccaccgctggtagcgggtggttttttgttgcaagcagcagattacgcgcagaa
aaaaaggatctcaagaagatcctttgatctttctacggggtctgacgctcagtggaacgaaaactcacgttaagggttttgg
tcatgagattatcaaaaaggatcttcacctagatccttttaaattaaaaatgaagttttaaataaatctaaagtatatatgagtaaa
cttggtctgacagttaccaatgcttaatacagtgaggcacctatctcagcgatctgtctatttcgttcatccatagttgcctgactcc
5 ccgtcgtgtagataactacgatacgggagggcttaccatctggccccagtgctgcaatgataccgcgagacccacgctcac
cgggtccagatttatcagcaataaaccagccagccggaagggccgagcgcagaagtggctcctgcaactttatccgcctcca
tccagctctattaattgttgcgggaagctagagtaagtagttccagtttaatagtttgcgaacgttggttccattgctacagg
catcgtggtgtcacgctcgtcgttgggtatggcttcattcagctccggttcccaacgatcaaggcgagttacatgatccccat
gttggtgcaaaaagcggtagctccttcggctcccgatcgtgtcagaagtaagtggccgcagtggtatcactcatggttatg
10 gcagcactgcataattctcttactgcatgccatccgtaagatgctttctgtgactggtgagtactcaaccaagtcattctgaga
atagtgatgacggcgaccgagtgctcttggccggcgtcaatacgggataataccgcgccacatagcagaactttaaaagtg
ctcatcattggaaaacgttctcggggcgaaaactcicaaggaacttaccgctgtgagatccagttcgatgtaacccactcgt
gcaccaactgatcttcagcatctttactttaccagcggttctgggtgagcaaaaacagggaaggcaaaatgccgcaaaaaa
gggaataaggcgacacggaaatgttgaatactcatactcttcttttcaatagctccaaggcaaaaatgactactcagac
15 cgacattcattcgttattgattttaaataacgataaacggaatggttacttgaatgatttcatttatgatcattgttactaattacc
taaataaggattttatatggaattggaagaataagggaatttcagatgctgaaaaaggcgaggagggtactaatcattcaag
cccatttctgccagtaattgcttcataagcttcaatatacttttcttactcttgatagcaatttctgcatccaaggctacgccctctt
tgccattcaatccgttggccgtcaaccaatctctgagaactgcttatcgtaactctcttgcgatttaccacttggtaagtctttt
gattccaaaatctagaagaatctggagttaaaactcatctactagtaccaattcattgtttcgtccagtcgaattcgaattcgt
20 atcagcaataatgatccccctcaaaaggcggaagtttttgcagcagaatacaactcgaccgccttgacagcgaccttctcac
aaatgtctttacctacaatctcagcagcttgttcaatagagatgttttcacgtgttcaccctgttcagcttctgttgaaggtgtgaa
aatcgggagttgaaaggcgctcgtctctgaagggtctcgttttcaaccttgactccatggacagttttgagttctgtactcttcc
catgcacttccagtgatgtaacctctgacaatggcttccaaaggatcagctctgtgtttttactatcaaggatcgtccctcta
tgagatttgattttctcagacagtttgatggtagtaagcaaaagacttcttgcattagaagcaaccaaatgattctttatgta
25 ggggtgccaaaaatcaaacgagaaaactgagagctgagtcaaaatctttcccttatcagggaataaccgtttgtcataatcacatc
gtaagcgggagatacggtcagttgcgacgaacagcaagttgttctcatcgactgcataaatgtcttaacctttccttggcgatt
aaaggtagattccgtccagatcagtggtcacaatggacatacttggaggatacagcaaaagtgtgttgaagcgatgacac
atggaaaggaattttcagatttctagtagtataattggggcggtgaaagttcagatgtttaatgcttaatactcttatactcttc
aaagcggcccaagtgttctgccaaactgacttttttgaataatgaatcgttcaagtggagtatttaaaccatgattaaagtacgt

gatttggcactgggataaggtcgaaaaatccgtatfcataaacgattattggtaaaagttacaaaataccactaattacggag
aagcttagtaacagttatcatctcttggcgttaaacgttacaatttccattcgccattcaggctgcgcaactgttgggaaggg
cgatcgggtgcgggcccctctcgtattacgccaggggcctcgaggcacaacgaacgtctcacttaattcttctglactctgaaga
ggagtgggaataaccaagaaaaacatcaaacctgaatgattttccaaacccctaccacaagatattcatcagctgcgagat
5 aggctgatcaggagcaagctcgtacgagaagaaacaaaatgacaaaaaaatcctatactatataagttacaataaaaaa
gtatcaaaaatgaagcctgcatctcagggcaaatggcattctgacatcctcttgattagaatctagcaagaccggctctctcgt
aagtggccaaactgaactgagggaacagtcattgtctaaaggctaaaactcaatgatgatgatgatggtcgcacggcgctattc
agatcctctctcgtagatgagttttgttcgggccccaaaaataacttcccaatcttcagttatgtcaagcgatagggttaagaatggt
gatcttattatcattctcatatgacgatctagtcggcaatgaactttctcgcattttttgactttaactgtcttcaacgttatgt
10 tgtgtgttaaattattcttgcgataatgatttttcaatattggtatcctcaaagtattggtgtgaatgcctgtcaaattctcgggaa
tatgacttcgaacatttttaaggatattgttttgaaaaatgaactgagttttctacgtactcacctcttggtagccaaaagtittcca
tcatcaacataaagatctccaacggctcgtccctcggtatcaggggctataactattacatatggatcgttttcattaacattga
agatcttctatactatcttcatagtataatgtgaccgccttcaataaataatggaattttatccaatgggtcgagaaatattcttt
ctatcaaatcagtaccattgttataaaagagttaaagatgcgaattcatagaatataccgggtgggaaaaccatttccgtttct
15 gattgaccaggctccgtgacaggttgactaatagacctgaattactccagtaaaattggttatcgatatgatacaattcagcaa
attcagggtgttcaataaacattggattcattatcggaatccagtgacacttgattatgaaacatggtgtataaggtaggttag
caggaaatatctcaattggataatacactgactatcgacttcaaagggtcattgaataagtafaggttctctcttctgtgtctata
tgggctgtgtctctaaaaatgggtaccataagcccgttggtaccaacgtgcaatcaattcagggtgtaggatcctcagcaa
agccagctatgtcggctcctataaatggcataccagcaalgtgtttgacagaaccatagggaatggaaatctttaagtaatccc
20 aattggccacattgtcaccagtcattgtggcagcagttgagagccggcaaaaaagcccttgttagaaggaaaggac
gcttatcggatggtgaataaatcattttattgcgtcgttaagtgccttcatgcactgatagaccatatgttatggacggatcttt
cctcaatgtaattgtcgtgaatcaaatctttggagctgtggtctctgggccatcgaaaatcgaggctcgttcataatcctcaa
atgaataaattagttaaatcagccggcagatccataaaccttcgaaaaaggacttccaaatctttggccatattgcttatggt
atcaatccatatagaattacctggccagcaatgacctacatagtcatttccattgtatccttgactgctacattttcatttaattacc
25 ctgtcactgatttcataatcttctttaaatgaggatcgttagtagcacaagatttctaccaactttttaattggataacagcct
ttttgattgggaaggagtgctgcttccaagtaaaatattttgtcgttctgtatctccaagtccaaccaataaaatcgtaag
gaatcatatgagcatccatctgagagtcacatgtgagaacgtccatctcatcattataattccatctacattgatggtacccatata
gaggaaatgggctgtaaaaagggtctaccagtcnaatcggtaaattgtcaatgatagttggaatactggccccaggggaca
tgactacatctatgacaccatttctggagatccaatgagtcacgttttttttactggtgtcactttatgtctaccaagtgtca

gctgcattgacccaaaagatagatgtggacgaagatgaaaacatgaatgggatcgaaccgtacattggttggctggtagc
 atgtgtactcaaagacatcaacgttgaaaagcctgtagggttccttccacctgaagtgtccatcagccttagcgcacgtcgca
 tgttccggtataccgtagacattagtagaaccatgaagagaaaactagcgcgaaccgattcaggcccaaaaggcatagag
 tcatgcttgaatacaagaaattgtccttaaacatgttgaaagtgtttcttctggcagcacgtgtgcgaagtttctcttagtgc
 5 tatgatgttcaatgttcaggaaatttgcctcgtttacaataagtttcagcgcattttgccagtaaaactttcaattgaaaaggttcagc
 aaagattttacggatacatcaccgttccgaagatgaaatgtgtctgcagtggagttgaaaagtacaaaaatgaagatatttc
 gaccagaatgagttcacagtttgttttgccttaaggagtggaattgtggaatactggctcctgttcgcctcctcttgaatttctgt
 cgaatgcgtacttccagggtctcattgaaccgttgtgaagagatcaacaaaccgctgtgttgggtggcatttctcttctcattt
 atagtgaaccttactgagtgatcctgtaaaaagagagagagaatgggaactgaacggctatatcatcgcctccaatcttg
 10 gtatagtttaattatggttagcatgaagcacattctctaaaggatcgtgtgcaatagactcggcgtccatttgaatagcagtg
 atgagatttggcaatattttctgcataaacctgttctatggcaaaaccagattgcgcacacttcttaatatagatagtcggtaa
 acgcatgcgaaaaagcggtaagaagaccaattggcatacgaccatttcaaaaggacatctcagggtaccgatccgag
 acggccggctgggccacgtgaattcttttctcagttgatttgttggggatttagtaagtcgtaaaacttcgttaaaaaagatca
 atgtagtcaatacagttgatccgaaatagaaggaagaggttgcattgtgaagaacaatgagttaaaagcccgttttaagac
 15 aatattctttagtgcgtgatcagaaaaggacaataagggaatttgggtgcttctttataccaataatcgtctcctcatcgcttaattt
 ctccccatctcaaccggtaggtaggacgttctgtaatctgttcacataaaaggggttttactccgagacaaaaatttat
 gcgacaaaaatagcctatcttggaaggatgtcttatcaactgcattgttgcaaggagaagcaaggacaactcaacatgg
 gtaaaaattcaaaaccaaccaattggaactcccaactgtccactagglagctgacagctgtcacttttctgttctgttctgt
 ctcttctgcttaactgtcctcttaccgcttcttccaaaaaattcgtgggaaatgtgccccatcagagtcctaatgacccat
 20 gaataaagtttctgtactgtttaagacgatgaattgcaacgataatccgagcagtttacggggtacatcacgtgctttgcatat
 gatctcggagtcggatcagttccggatgtgatgtattacccatagtttcaactctaatgcagccgccaagtgcatacacc
 ctccatcaatctatgcttaagttttaccatcgttgggtgggtgatgatgactcgttagtctctgtgttcgatattaactttgtaa
 ggatcgccttggatggaaaattgaggggttgaacctgaatttcaggctacttacattggacttttgagaaggctggacgg
 ttgatgaagagggtgggtgcagaggaatgaaaaaaatttagttgagaggactgcttgaattttaggaatggagtccttt
 25 aagctgacaaaactcaaggatggggtatttcatgtagcttttcatgccttcgacaagctaaagggaagtaattgattctggat
 aatggatatttgatctgcttagcagatgtcaaaagtctactagtatagctcgtatctcgtagccttcaattggcgatctta
 ctgaaagtgttatatttttagctgacgagacgaagaacgagagagtattgacacattcagaggtaagacaatatgtcgtattat
 caaaataagtatcgaacctctattaggagcctactggctcaaatgtgcaaccttagtggtgattgtctctgcttctgacacaaat
 ctgtcgtgttgagagtgccgatgtatgatttttagtaaatgttttcagaaaaggcgtaagtaaaataaccagtaagtaataaat

aacgtaaaagtgtttgaatcataaaagaatcaagatagagggtcaaagcatagataatcccccgtagaagatggaccggtc
atatggctgaaaaaagatctgatctcatg

SEQ ID NO: 18 pGAPZAGLSII

5 tcgagatggctctttgaaatggctcgtatgccaattggctcttttaccgcttttcgcatgcgtttaccgactatctattaagaa
gtgtgcgcaatctgggttttggcatagaaacagggtttatgcagaaaatttggcaaatctcatcactgctattacaaagtga
cgccgagtctattgcacacgatccttagagaatgtgcttcatgctaccataattaaaactataccaagattggaggcgatga
tatagccgttcagttccattctctctctctttttacaggatcacicagtaagggtcactataaatgagaaagagagaatgccaa
ccaacagcagcggttgtgatctcttcacaacgggtcaatgagacctggaagtacgcattcgacaagaaattcaagagga
10 ggcgaacaggaccagtattccacaattccacttccctaagcaaaaactgtgaactcattctggtcgaaaaatctctcatit
ttgtcactttcaactccactgcagacacattcatcttgcgaacgggtgatgtatccgtagaaatcttctgtaacctttcaattg
aaagttaactggcaaatgcgctgaaacttattgtaaacgagcaaaatttctgaacattgaacatcatagaactaagcagga
aaacttcgcacacgtgctgccagaagaacaacttcaacatgtttaaggacaatttctgtattcaagcatgactctatgcctt
tggggcctgaatcggttgcgctagatttctcttcatgggttctactaatgtctacgggtataccggaaatgcgacgtcgtctaa
15 gctgatggacacttcaggtggaagggaacctacaggcttttcaacgttgaatgtctttgagtacaacatcggtaccagccaac
caatgtacgggtcgtatccattcatgttttcatcttctccacatctatcttttgggtcaatgcagctgacacttgggtagacataa
agtatgacaccagtaaaaaataaacgatgactcattggatctccgaaaatgggtcatagatgtatcatgtccctggggcca
gatattccaactatcattgacaaatttaccgatttgactggtagaccttttaccgcccatttctctatagggtaccatcaatgta
gatggaattataatgatgatgagatggacgttctcacagtggactctcagatggatgctcatatgattccttacgttttatttgggtg
20 acttggagtatacgaacgacaaaaaatattttacttgggaagcagcaccttcccaatccaaaaggctgttatccaaattaa
aaagtgggtagaaatcttctgtactaatcgtatccatcttaagaaagattatgaaatcagtgacagggttaattaatgaaat
gtagcagtcaaggatcacaatggaaatgactatgtaggtcattgctgccaggtaattctatatggattgataccataagcaaaa
tatggccaaaagatttgaagtccttttgaacgggttatggtatctgccggctgatitaaactaatttattcatttgaatgatatga
acgagccttcgatttctgatggccagagaccagctccaaaagatttgattcacgacaattacattagggaagatccgtc
25 cataacatafatggctcatcagtgcataagctacttacgcacgaataaaatcgatttattcaccatccgataagcgtcctttcct
tctaacaagggttttttgcggctctcaacgtactgctgccacatggactgggtgacaatgtggccaattgggtacttaaaag
atttccattcctatggttctgtcaacaacattgctggatgccatttataggagccgacatagctggcttctgaggatccttac
acctgaattgattgcacgttgggtaccaagcgggttatgggtaccatttttagagcacacgcccataatagaccaagagaa
gagaaccatacttattcaatgaaccttgaagtcgatagtagctgatattatccaattgagatatttctgctacctacctatataca

ccatgtttcataaatcaagtgtcactggatttccgataatgaatccaatgtttattgaacacctgaatttctgaattgtatcatat
 cgataaccaattttactggagtaaitcaggtctattagtaaacctgtcacggagccgtgtcaatcagaaacggaaatggtttc
 ccacccgggtatattctatgaattcgcacatcttttataaacaatgggtactgatttgatagaaaagaataattctgcacat
 tggataaaaattccattatttgaaggcgggtcacattatcactatgaagataagtatagaagatcttcaatgttaataaaaaac
 5 gatccatattgaatagttatagcccctgataccgaggagcagccgtggagatctttatgttgatgatggagaaactttggct
 accaaagagggtgagtagtagaaactcagttcattttcgaaaacaataaccttaaaaaatgttcgaagtcataatcccgagaattt
 gacaggcattcaccacaatactttgaggaataccaatattgaaaaatcattatcgcaaagaataattacaacacaacataac
 gttgaaagacagtattaaagtcaaaaaaatggcgaagaaagttcattgccgactagatcgtcatatgagaatgataataaga
 tcaccattctaacctatcgcttgacataactgaagattgggaagtattttggggccgaacaaaaactcatctcagaagagg
 10 atctgaatagcgcctcgacatcatcatcatcatcattgagtttagccttagacatgactgttcctcagttcaagttgggcact
 tacgagaagaccggtcttctagattctaatcaagaggatgtcagaatgccatttgcctgagagatgcaggcttcattttgata
 cttttttttgtaacctatatagtataggattttttgtcattttgttcttctcgtacgagcttgctcctgatcagcctatctcgagct
 gatgaatatcttgggtagggtttgggaaaatcattcgagtttgatgttttcttggtatttccactcctctcagagtacagaa
 gattaagtgagaccttcgttgcggatccccacacacacatagcttcaaatgtttctactcctttttactcttccagattttctc
 15 ggactccgcgcacgcgtaccacttcaaacacccaagcacagcactataaattttcccttttctcctctagggtgtcgtta
 attaccgactactaaagggttggaaaagaaaaagagaccgctcgtttcttctcgtcgaanaaggaataaaaaattttatc
 acgtttcttttctgaaatttttttttagttttttcttctcagtgacctccattgatatttaagtaataaacggcttcaatttctcaa
 gtttcagtttcaattttctgttctattacaacttttttacttctgttcattagaaagaaagcatagcaatctaatcaaggcggtgt
 tgacaattaatcatcgccatagtatatcgcatagatataacgacaaggtaggaactaaacctggccaagttgaccagtg
 20 ccgttccggtgtcaccgcgcgcgacgtcggcgagcgggtcagttctggaccgaccggctcgggttctccgggacttc
 gtggaggacgacttcggcggtgtgtcgggacgacgtgacctgttcacgcgcggtccaggaccaggtgtgtccgg
 aacacacctggcctgggtgtgggtgcgcggcctggacgagctgtacgcgagtggtcggaggtcgtgtccacgaacttc
 cgggacgctcggggccggccatgaccgagatcggcgagcagccgtgggggaggagttcgccctgcgcgacccgg
 ccggcaactgcgtgcacttcgtggccgaggagcaggactgacacgtccgacggcgccacgggtcccaggcctcggg
 25 gatccgtcccccttttcttctgataatcatgtaattagttatgtcacgcttacattcacgccctccccacatccgctctaacc
 gaaaagggaaggagttagacaacctgaagtcaggtccctattttttatagttatgttagtattaagaagcttatttatattca
 aattttctttttctgtacagacgcgtgtacgcatgtaacattatactgaaaaccttgcctgagaagggtttgggacgctcgaa
 ggctttaatgtgaagctggagaccaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccgcttg
 ctggcgttttccataggtccgccccctgacgagcatcacaataatcgacgctcaagtcagagggtggcgaaacccgaca

ggactataaagataccaggcggttccccctggaagctccctcgtgcgctctcctgttccgacctgccgcttacgggatacct
 gtccgcctttctcccttcgggaagcgtggcgctttctcaatgctcacgctgtaggtatctcagttcggtgtaggtcgctcctcc
 aagctgggctgtgtgcacgaacccccgttcagcccagccgctgcgccttatccggtaactatcgtcttgagccaacccgg
 taagacacgactatcgccactggcagcagccactggtaacaggattagcagagcgaggatgtagggggtgtacagag
 5 tcttgaaagtggtggcctaactacggctacactagaaggacagtattggatctgcgctctgctgaagccagttacctcggg
 aaaagagttggtagctctgatccggcaacaacaccaccgctggtagcggtggtttttgttgcaagcagcagattacgcg
 cagaaaaaaggatctcaagaagatccttgatctttctacggggtctgacgctcagtggaacgaaaactcacgttaaggg
 attttggtcatgcatgagatcagatctttttagaaatgcttgggtgctcctcgtccaatcaggtagccatctctgaaatatctgg
 ctccgttgcaactcgaacgacctgctggcaacgtaaaatctccgggtaaaactaaatgtggagtaaatgaaccagaaa
 10 cgtctctccctctctctccttccaccgcccgttaccgtccctaggaattttactctgctggagagcttctctacggccccctt
 gcagcaatgctcttccagcattacgttgcgggtaaacggaggtcgtgtacccgacctagcagcccagggtggaag
 tcccgccgctcgtggcaataatagcggggcgacgcatgcatgagattattgaaaccaccagaatcgaalataaaaaggc
 gaacaccttcccaatttgggttctcctgacccaagactttaaatttatttgcctatttcaatcaattgaacaactattc
 gaaacgagggaattcacgtggccagccggccgtctcggtacgggtacc

15

SEQ ID NO: 19 pGAPADE1glsII

tcgaccggctgcattaatgaatcgccaacgcgcggggagagcgcggttgcgtattgggcgctcttccgcttctcgtcac
 tgactcgtcgcgcgtcggtcgttcggctgcggcgagcgggtatcagctcactcaaaggcggtataacggttaccacagaatca
 ggggataacgcaggaagaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaggccgcttgcgtggc
 20 gttttccataggctccgccccctgacgagcatcacaataatcgacgctcaagtcagaggtggcgaaacccgacaggact
 ataaagataccaggcggttccccctggaagctccctcgtgcgctctcctgttccgacctgccgcttacgggatacctgtccg
 cctttctcccttgggaagcgtggcgctttctcatagctcacgctgtaggtatctcagttcggtgtaggtcgttcgctccaagct
 gggctgtgtgcacgaacccccgttcagcccagccgctgcgccttatccggtaactatcgtcttgagccaacccggtgaaga
 cagcacttatcgccactggcagcagccactggtaacaggattagcagagcgaggatgtagggcggtgtacagagttcttg
 25 aagtggtagcctaactacggctacactagaaggacagtattggatctgcgctctgctgaagccagttaccttcggaaaaag
 agttggttagctcttgatccggcaacaacaccaccgctggtagcggtggtttttgttgcaagcagcagattacgcgcagaa
 aaaaaggatctcaagaagatccttgatctttctacggggtctgacgctcagtggaacgaaaactcacgttaagggttttgg
 tcatgagattatcaaaaaggatcttcacctagatccttttaataaaaaatgaagtttaaatcaatctaaagtatatatgagtaaa
 ctgtgtctgacagttaccaatgcttaacagtggagccatctcagcgatctgtctatttcgtcatccatagttgcctgactcc

ccgtcgtgtagataactacgatacgggagggcttaccatctggccccagtgctgcaatgataccgcgagacccacgctcac
 cggctccagatttatcagcaataaaccagccagccggaagggccgagcgagaagtggctcgaactttatccgcccca
 tccagttctattaattgttgcgggaagctagagtaagtagttccagtttaatagtttgcgcaacgttgttgcattgtacagg
 catcgtgggtgcacgctcgtcgtttggtatggcttcattcagctccgggtcccaacgatcaaggcgagttacatgatccccat
 5 gttgtgcaaaaaagcgggttagctccttcggctcccgatcgtgtgcagaagtaagtggccgcagtggtatcactcatggttatg
 gcagcactgcataattctcttactgtcatgccatccgtaagatgcttttctgtactggtgagtactcaaccaagtcattctgaga
 atagtgatgcggcgaccgagttgctcttgcggcggtcaatacgggataataccgcgccacatagcagaactttaaagtg
 ctcatcattggaaaacgttcttcggggcgaaaactctcaaggatcttaccgctgttgagatccagttcgatgaaccactcgt
 gcaccaactgatcttcagcatctttactttaccagcgtttctgggtgagcaaaaacaggaaggcaaaatgccgcaaaaaa
 10 gggaataaggcgacacggaaatgttgaatactacalcitcttcttttcaatagctccaaggcaaaaatgactactcagac
 cgacattcattcgttattgattttaaatacagataaacggaatggttacttgaatgattcactttatgatcattgttactaattacc
 taaataggattttatatggaattggaagaataagggaatctcagatgtctgaaaaaggcgaggagggtactaatcattcaag
 cccatttcttgcagtaattgcttcataagcttcaatatacttttcttactcttgatagcaatttctgcatccatggctacgccctctt
 tgcattcaatccgttggcgtcaaccaatctctgagaaactgcttatcgaactctcttgcgatttaccacttggttaagtcttt
 15 gattccaaaatctagaagaatctggagttaaaacttcatctactagtaaccaattcattgtttcgtccagttcaaaatcgaatttgc
 atcagcaataatgatccccctcaaaaggcggaagtttttgcagcagaatacaactcgaccgccttgacagcgaccttctcac
 aaatgtctttacctaacaatcagcagcttgttcaatagagatgtttcatcgtgttcacctgttcagcttctgtgaaggtgtgaa
 aatcggagttgaaaggcgtcgtctcttgaaggtctcttcaaccttgactccatggacagtitttgagtcttctgactctttc
 catgcacttccagtgatgtaacctctgacaatggcttcaaaaggtatcagctgtgtcttttactatcaaggatcgtccctcta
 20 tgagatttgattttctcagacagttttgatgtagtaagcaaaagacttcttgcattagaagcaaccaaatgaitctttatgta
 ggggtgccaaaaaatcaaacagaaaaactgagagctgagtcaaaactttcccttatcaggaataccgtttgcataatcacatc
 gtaagcggagatacggtcagttgcgacgaacgaagttgttctcatcgactgcataaatgtctctaactttcccttggcgatt
 aaaggtaggattccgtccagatcagtttcacaatggacatacttgaaggatacagcaaaagtgtgttgaagcgatgacac
 atggaaaaggaaattttcagagttcctagagtagtatattggggcggtgaaagttcagatgtttaatgcttaatactcttatactctc
 25 aaagcgcccaagtgtttctgccaacctgactttttctgaataatgaatcgttcaagtggagtatttaaccatgattaaagtacgt
 gatttggcactggataaggctgaaaaatatccgtattcataaacgattattggtaaaagttacaaaataaccactaattacggag
 aagcttagtaacagttatcatctcttggctgattaacgcttacaatttccattcgcattcaggctgcgcaactgttgggaaggg
 cgatcgggtgcgggcctcttgcattacgccagggcctcgaggcacaacgaacgtctcacttaactcttctgactctgaaga
 ggagtggggaataccaagaaaaacatcaaacgaatgattttccaaaccctaccacaagatattcatcagctgcgagat

aggctgatcaggagcaagctcgtacgagaagaacaaaatgacaaaaaaatcctatactataggttacaataaaaa
gtatcaaaaatgaagcctgcatctctcaggcaaatggcattctgacatcctcttgattagaatctagcaagaccggctctctcgt
aagtgcccaacttgaactgagggaacagtcagtcctaaagctaaaactcaatgatgatgatgatggcgacggcgctattc
agatcctctctgagatgagttttgttcggggcccaaaaataacttcccaatcttcagttatgtcaagcgatagggttaagaatggg
5 gatcttattatcattctcatatgacgatctagtcggcaatgaacttcttcgccattttttgactttaatactgtctttcaacggttatgt
tgtgttgtaaatattctttgcgataatgatttttcaatattgggtattcccaaagtattgtggtgaatgccgtgcaaatctcgggaa
tatgacttcgaacatttttaagggtattgtttcgaaaatgaactgagttctacgtactcacctcttggtagccaaaagtttctcca
tcatacaataaagatctccaacggctcgtccctcggtatcaggggctataactattacatatggatcgttttcattaacattga
agatcttctatacttattcttcagtgataatgtgaccgccttcaataaataatggaattttatccaatgggtcgagaaatattcttt
10 ctatcaaatcagtaccattgtttataaaagagtgtaaagatgcgaattcatagaataaccgggtgggaaaaccatttccgtttct
gattgaccaggctccgtgacagggttgactaatagacctgaattactccagtaaaatgggtatcgatatgatacaattcagcaa
attcagggtgttcaataaacattggattcattatcggaatccagtgacacttgatttatgaaacatgggtgtataaggtaggtag
caggaaatatctcaattggataatatcacgtactatcgacttcaaagggtcattgaataagtatgggtctctctcttgggtgtctata
tgggctgtgtcttaaaaaatgggtaccataagcccgttgggtaccaacgtgcaatcaattcagggtgtaggatcctcagcaa
15 agccagctatgtcggctccataaatggcalaccagcaatgttgttgacagaaccataggaaatcfttaagtaatccc
aattggccacattgtcaccagtcattgtggcagcagtcaggttgagagccggcaaaaaagcccttggtagaaggaaaggac
gcttatcggatgggtgaataaatcattttattgcgtcgttaagtagcttcatgcactgatagaccatatgttatggacggatcttl
cctcaatgtaattgtcgtgaatcaaatcttttgagctgtgtgtctctgggccatcgaaaatcgaggctcgttcatatcattccaa
atgaataaattagttaaatcagccggcagatccataaacggttcgaaaaaggacttccaaatctttggccatatttgcttatgtt
20 atcaatccatatagaattacctggccagcaatgacctacatagtcatttccattgtgaccttgactgtacattttcattaattacc
ctgtcactgatttcataatctttttaaattgaggatcgatttagtcagacaagatttctaccaacttttttaatttgataacagcct
ttttggattgggaaaggagtgtgttccaagtaaaatattttgtcgttcgtatactccaagtccaaccaataaaatcgtaag
gaatcatatgagcatccatctgagagtcacatgtgagaacgtccatctcatcattataattccatctacattgatggtagccctata
gaggaaatgggctgtaaaaagggtctaccagtcacaaatcggttaatttgcattgatagttggaatatctggccccagggaca
25 tgactacatctatgacaccattttcgagatccaatgagtcacgttttttttacttggtgtcactttatgtctaccaagtgtca
gctgcattgacccaaaagatagatgtggacgaagatgaaaacatgaatgggatcgaaccgtacattggttgctggtaccg
atgtgtactcaagacatcaacgttgaaaagcctgtagggttcctttccacctgaagtgtccatcagccttagcgacgtcgca
tgttccggtataccgtagacatttagtagaacccatgaaagagaaatctagcgcaaccgattcaggcccaaaaggcatagag
tcatgctttgaatacaagaatgtccttaacatgttgaaagttgtttctctggcagcacgtgtgcgaagtttctgtcttagttc

tatgatgttcaatgttcaggaaattttgctcgtttacaataagtttcagcgcatittgccagtaaactttcaaltgaaaagggttcagc
 aaagattttcaccgatacatcaccgtttcgaagatgaaatgtgtcgcagtgagggttggaaagtacaaaaatgaagatattttc
 gaccagaatgagttcacagtttgttttgcttaaggagtggaattgtggaatactggctcgttcgcctcctcttgaatttcttgt
 cgaatgctgacttcagggtctcattgaaccgttgtgaagagatcaacaaaccgctgctgttggttggaacttctcttctcattt
 5 atagtgaaccttactgagtgatcctgtaaaaagagagagagaatgggaactgaacggctatatcatcgccctccaatcttg
 gtatagtttaattatgtagcatgaagcacatttctaaggatcgtgtgcaatagactggcgtccactttglaatagcagtg
 atgagatttggcaatatcttgcataaacctgttctatggcaaaaccagattgcgcacacttcttaatatagatgctgtaa
 acgcatgcgaaaaagcggtaagaagaccaattggcatagcagccatttcaaaaggaccatctcgaggtaccgatccgag
 acggccggcgtggccacgtgaattcctcgttcgaaatagttgttcaattgattgaaataggggacaaataaattaaattaaagt
 10 ctttgggtcaggagaaacaaaattgggaagggtgttcgcctttatattcgattctggtggttccaataatctcatgacatgcg
 tccgcccgctattattgccagcgacggccgggactttccatccctgggctgctaggtcgggtacacgacctccgtttaccoc
 gcaacgtaatgctgggaagagcattgtgcaaggggcgtagaagaagcttccagcagagtaaaatttctaggggacg
 gtaacggcggtgggaaggagagagaagggaagagacgtttctgttccattactccacatttaagtttaccgggagaattt
 tacgttgccagcaggtcgttcggagttgcaacggagccagataittcagagatggctacctgattggacgaggacaccaag
 15 acatttctacaaaaagatctgatctca

SEQ ID NO: 20 pPICZAGLSII

cgaacaaaaactatctcagaagaggaatcgaatagcggcgtcgaccatcatcatcatcatcattgagttgtagccttagac
 atgactgttctcagttcaagttgggcacttacgagaagaccggcttctgtagatttctaataagaggatgtcagaatgccattt
 20 gcctgagagatgcaggcttcattttgatactttttatttgaacctatatagtataggattttttgtcattttgttcttctcgtacga
 gcttctcctgatcagccatctcgcagctgatgaatacttctgtgtaggggtttgggaaatcattcgagtttgatgttttcttg
 gtatttcccactcctcttcagagtacagaagattaagtgaaccttctgttgcggatccccacacaccatagcttcaaaatg
 ttttactccttttttactcttcagattttctcgactccgcgcacgcgtaccacttcaaaacaccaagcacagcactataa
 attttcccttcttctcctciagggtgtcgtaattaccggtactaaagggttggaaaagaaaaagagaccgctcgtttctttt
 25 ttctgcgaaaaaggcaataaaaaattttatcacgtttcttttcttgaatttttttttagtttttcttcttcagtacctccattgat
 ttaagttaataaacgggtcttcaatttctcaagtttcagttcatttttctgttctattacaacttttttacttctgttcattagaaagaa
 gcatagcaatctaataaggggcggtgttgacaattaatcatcgccatagatatacgcatagataatacgaaggtgag
 gaactaaaccatggccaagttgaccagtgccgttccgggtgctcaccgcgcgcgacgtcggcgagcgggtcaggttctgga
 ccgaccgggtcgggttctccgggacttctgtggaggacgacttccgggtgtgttcgggacgacgtgacctgttcatca

gcgcgggtccaggaccagggtgggtgccggacaacacctggcctgggtgtgggtgcgcggcctggagagctgtacgccg
agtggtcggaggtcgtgtccacgaacttccgggacgcctccggggccatgaccgagatcggcgagcagccgtggg
ggcgggagttccctgcgcgacctggccggcaactgcgtgcacttcgtggccgaggagcaggactgacacgtccgac
ggcggcccacgggtcccaggcctcgagatccgtccccccttctgtcgatcatgtaattagttatgtcacgcttacatt
5 cagccctccccacatccgctctaaccgaaaaggaggagtagacaacctgaagtctaggctccattatattttatagt
tatgttagtattaagaacgttatttatattcaaattttctttttctgtacagacgcgtgtacgcatgtaactataactgaaaacc
ttccttgagaaggtttgggacgcctcgaaggcttlaatttgcaagctggagaccaacatgtgagcaaaaggccagcaaaag
gccagggaaccgtaaaaaggccgcgttgcgtggcgttttccataggctccgccccctgacgagcatcacaaaaatcgacgc
tcaagttagagggtggcgaacccgacaggactataaagataccaggcgttccccctggaagctccctcgtgcgtctcct
10 gttccgacctgccgcttaccggatacctgtccgctttctcccttcgggaagcgtggcgttttctaatgctcacgctgtagg
tatctcagttcgggtgtaggctgttcgctccaagctgggctgtgtgcacgaacccccgttcagcccagccgtcgcgcttate
cggtaactatcgtcttgagccaacccgtaagacacgaactatcgccactggcagcagccactggtaacaggattagcag
agcgagggtatgtaggcgggtgtacagagttctgaagtggtggcctaactacggctacactagaaggacagattttggtatct
gcgctctgctgaagccagttaccttcggaaaaagagttggtagctcttgatccggcaacaaaccaccgctggtagcgggtg
15 gttttttgttgcaagcagcagattacgcgcagaaaaaaggatcicaagaagatcctttgatcttttctacggggtctgacgc
tcagtggaaacgaaaactcacgttaagggttttggctatgagatcagatcfaacatccaagacgaaaggttgatgaaacct
tttgcacccgacatccacaggttcattctcacataagtgcacaaacgcaacaggaggggatacactagcagcagaccg
ttgcaaacgcaggacctccactcctcttctcctcaacacccacttttgccatcgaaaaaccagcccagttattggcgttgattg
gagctcgtcattccaattcctctattaggctactaacacacatgactttatagcctgtctatcctggccccctggcgagggttc
20 atgtttgtttatttccgaatgcaacaagctccgattacacccgaacatcactccagatgagggtcttctgagttgtgggtcaa
atagttcatgttccccaaatggcccaaaactgacagtttaaacgtgtcttggaacctaatatgacaaaagcgtgatctcatcc
aagatgaactaagtttgggtcgttgaaatgctaacggcagttggcctcaaaaagaacttccaaaagtcggcataccgtttgtct
tgtttggtattgattgacgaatgctcaaaaataatctcattaatgcttagcgcagtcctctatcgtcttgaaccccggtgcacct
gtgccgaaacgcaaatggggaaacacccgcttttggatgattatgcattgtctccacattgtatgcttcaagattctggtgg
25 gaatactgctgatagcctaacgttcatgatcaaaatttaactgttctaaccctacttgacagcaatatataaacagaaggaag
ctgccctgtcttaaaccttttttatacatcattattagcttacttcataattgcgactggttccaattgacaagctttgatttaacg
actttaacgacaacttgagaagatcaaaaaacaactaattatcgaaacgaggaattcacgtggcccagccggccgtctcg
gatcggtagctcgagatggctcctttgaaatggctcgtatgccaattggctctttaccgcttttgcacgtgttaccgactat
ctattaaagaagtggtgcgaatctgggtttgccatagaacagggttatgcagaaaatattgccaatctcatcactgctatt

acaaagtggacgccgagtcattgcacacgatcccttagagaatgtgcttcattgctaccataaataaaactataccaagattgg
 agggcgatgatatagccgttcagttccattctctctctttttacaggatcactcagtaagggtcactataaatgagaaagag
 agaatgccaaccaacagcagcgggttgatctcttcacaacgggtcaatgagacctggaagtacgcattcgacaagaaatt
 tcaagaggaggcggaacaggaccagtattccacaattccacttccctaagcaaaaacaaactgtgaactcattctggtcgaaa
 5 atatcttcattttgtcactttcaaacccactgcagacacattcatcttcgaaacgggtgatgtatccgtagaaatcttggcgaac
 cttttcaattgaaagttaactggcaaatgcgtgaaactattgtaaacgagcaaaatttccgaactgaacatcatagaact
 aagcaggaaaaacttcgcacacgtgctgccagaagaaacaactttcaacatgtttaaggacaatttctgtattcaagcatga
 ctctatgcctttggggccgaatcggttcgctagatttctcttcatgggttctactaatgtctacgggtataccgggaacatgcga
 cgtcgctaaggctgatggacacttcagggtgaaaggaaacctacaggcttttcaacgttgatgtctttgagtacaacatcggt
 10 accagccaaccaatgtacgggttcgatccattcatgtttcatcttcgtccacatctatctttgggtcaatgcagctgacacttg
 ggtagacataaagtatgacaccagtaaaaaataaacgatgactcattggatctccgaaatgggtgcatagatgtatgcatgt
 ccctggggccagatattccaactatcattgacaaattaccgattgactggtagaccttttaccgccatttctctataggg
 taccatcaatgtataggaattataatgatgagatggacgttctcacagtggactctcagatggatgctcatatgattccttacg
 attttatttgggtgacttgaggtatacgaacgacaaaaataatttacttggaaagcagcactccttcccaatccaaaaaggctg
 15 ttatccaaataaaaaagttgggtagaatctgtcgtactaatgatcctcatttaagaaagattatgaaatcagtgacagggt
 aattaatgaaaatgtagcagtcaggatcacaatggaaatgactatgtaggtcattgctggccaggtaattctatatggattgat
 accataagcaaatatggccaaaagatttggaaagtccttttgaacgggttatggatctgccggctgatttaactaatttattcatt
 tggaatgatatgaacgagcccttcgattttcgatggcccagagaccacagctccaaaagatttgattcacgacaattacattga
 ggaaagatccgtccataacatfatgggtatcagtgcatgaagctacttacgacgcaataaaatcgatttattcaccatccgat
 20 aagcgtcctttccttaacaagggttttttggccggtctcaacgtactgctgccacatggactgggtgacaatgtggccaatt
 gggattacttaagatttcattcctatgggtctgtcaacaacattgctggatgccatttataggagccgacatagctggcttt
 gctgaggatcctacacctgaattgattgcacgttggtaaccaagcgggcttatggtaccatttttagagcacacgcccataa
 gacaccaagagaagagaaccatactattcaatgaaccttgaagtcgatagtacgtgatattatccaattgagatattcctgc
 tacttacctatataccatgtttcataatcaagtgtcactggatttccgataatgaatccaatgtttattgaacacctgaatttgc
 25 tgaattgtatcatacgaataccaatttactggagtaattcaggtctattagcaaacctgtcacggagcctggfcaatcagaaa
 cggaaatgggtttccaccgggtatattctatgaattcgcatctttacactctttataaacaatggtacttgattgatagaaaaga
 atatttctgcaccattggataaaattccatttattgaaggcgggtcacattatcactatgaagataagtatagaagatcttcaa
 tgtfaatgaaaaacgatccatatgtaatagttatagcccctgataccgaggagccggttgagatctttatgttgatgatg
 gagaacatttggctaccaaaagggtgagtacgtagaactcagttcattttcgaaaacaataccttaaaaaatgttcgaagtc

atattcccagaatttgacaggcattcaccacaatactttgaggaataccaatattgaaaaatcattatcgcaaagaataatt
acaacacaacataacgttgaaagacaglattaaagcaaaaaaatggcgaagaaagtcattgccgactagatcgcatat
gagaatgataataagatcaccattcttaacctatcgcttgacataactgaagattgggaagtatttttgggcc

5 **SEQ ID NO: 21** pPICADE1glsII

aaattcctcgtttcgaataattagttgttttgaatcttcaagttgctgtaaaagtcgttaaaatcaaaagctgtcaattggaac
cagtcgcaattatgaaagtaagctaataatgatgataaaaaaaagggttaagacagggcagcttccttctgttatattgct
gtcaagtaggggttagaacagttaaatttgaiccatgaacgttaggctatcagcaglattccaccagaatcttggaagcatac
aatgtggagacaatgcataatcatccaaaaagcgggtgttcccatcttgcgttcggcacagggtgcaccgggggttcagaag
10 cgatagagagactgcgctaagcattaatgagattattttgagcattcgtcaatcaatccaaacaagacaacgggatgccg
acttttgaagttcttttgaccaactggccgttagcatttcaacgaacaaactlgttcatcttgatgagatcacgctttgtc
atattagggtccaagacagcgtttaaactgtcagtttgggccatttgggaacatgaactatttgacccccactcagaag
ccctcatctggagtgatgttcgggtgtaatgcggagcttggtcattcggaaataaacaacatgaacctgccaggggggc
caggatagacaggctaataaagtcagtggttagtagcctaatagaaggaaattggaatgagcgagctccaatcaagcccaat
15 aactgggctggttttgcagtgcaaaagtgggtgttaggagaagaggagtgaggctcctgcgttgcaacggctgctgct
agtgtatccctcctgttgcgtttggcacttatgtgtgagaatggacctgtggatgtcggatggcaaaaagggttcattcaacct
ttcgtctttggatgttgcgaccggctgcattaatgaatcgcccaacgcgcggggagaggcggttgcgtattgggcgctctt
ccgcttcctgcctcactgactcgtcgcctcggctgttcggctgcggcgagcgggtatcagctcactcaaaaggcggtataatcg
gttatccacagaatcaggggataacgcaggaaagaacatgtgagcaaaaggccagcaaaaggccaggaaccgttaaaaa
20 ggccgcgttgctggcggttttccataggctcggccccctgacgagcatcacaaaaatcgacgctcaagtcagaggtggcg
aaaccgcagaggactataaagataaccaggcggttccccctggaagctccctcgicgcgtctcctgttcgacctgccgctt
accggatacctgtccgccttctccctcgggaagcgtggcgctttctcatagctcacgctgtaggtatctcagttcgggttag
gtcgttcgtccaagctgggtgtgtgcaagaacccccgttcagcccaccgctgcgccttatccggttaactatcgtcttga
gtccaaccggtaagacacgacttatcgccactggcagcagccactgtaacaggattagcagagcgaggtatgtagcg
25 gtgctacagagttcttgaagtgtggcctaactacggctacactagaaggacagtatttggtaictgcgtctgctgaagcca
gttaccttcggaagaggttggttagctcttgatccggcaaaacaaccaccgctgtagcgggtgttttttgttgcaagcag
cagattacgcgcagaaaaaaggatctcaagaagatccttgaatctttctacggggtctgacgctcagtggaacgaaaactc
acgttaagggaatttggatgagattatcaaaaaggatcttcacctagatccttttaattaaaaatgaagttttaaatcaatcta
aagtatatagtaaaacttggctgacagttaccaatgcttaacagtgaggcacctatctcagcgatctgtctatttcgttcac

catagttgcctgactccccgtcgtgtagataactacgatacgggagggttaccatctggccccagtgctgcaatgataccg
 cgagaccacgctcaccggctccagattatcagcaataaaccagccagccggaaggccgagcgcagaagtggctctg
 caactttatccgctccatccagctatattaattgttgcgggaagctagagtaagtagttccgccaagtttgccgaacgtt
 gttgccattgctacaggcatcgtgggtcacgcctcgttgggtatggcttcattcagctccgggtcccaacgaacaggcga
 5 gttacatgatcccccatgttgtcaaaaaagcgggttagctccttcggctcctccgatcgtgtcagaagtaagtggccgagtg
 ttatcactcatggttatggcagcactgcataattctcttactgtcatgccatccgtaagatgctttctgtgactggtagtactca
 accaagtcattctgagaatagtgtatcgggcgaccgagttgctcttggccggcgtcaatacgggataataccgcccacata
 gcagaactttaaaagtctcatcattggaaaacgttctcggggcgaaaactcicaagatctaccgctgttgagatccagtt
 cgatgtaaccacactcgtgcaccaactgatcttcagcatctttacttaccagcgtttctgggtgagcaaaaacaggaaggc
 10 aaaaatgccgcaaaaaagggaataaggcgacacggaaatgtgaatactcatacttcttcttttaaatagctccaaggcaac
 aaattgactactcagaccgacattcaltcgttatgtatttaataacgataaacggaatggtacttgaatgatttactttatga
 tcattgtttactaattacctaataaggattttatggaattggaagaataagggaattcagatgtctgaaaaaggcgaggag
 ggtactaatcattcaagccatttcttgcagtaattgcttcataagcttcaataacttttcttactcttgatagcaatttctgcac
 catggctacgccctcttgcattcaatccgttggccgtaaccaatctctgagaaactgctatcgaactctcttgcgatttac
 15 ccacttggtaagcttttgattccaaaatctagaagaatctggagttaaaacttcatctactagtagccaattcattgtttcgtccag
 tccaaatcgaattcgtatcagcaataatgatcccttcaaaaggcggaagtttttgcagcagaatacaactcgaccgccttg
 acagcgaccttctcacaatgtctttacataatctcagcagcttgcataagagatgtttcatcgtgttcacctgttcagct
 ttcgttgaagggtgaaatcgagttggaaggcgctcgtctcttgaagggtctcgtttcaaccttgactccatggacagttt
 tgagttcttgactcttccatgcacttccagtgaatgaacctctgacaatggcttcaaggtatcagctctgtgtttttactatca
 20 aggatcgtccctctaattgagatttgtattttcttcagacagttttagtgtagtaagcaaagacttcttgcattagaagcaa
 ccaaatgattctttatgtagggtgccaaaaaatcaaacgaaaaactgagagctgagtcataatcttcccttaccaggaatac
 cgtttgcataatcacatcgttaagcggagatacggtcagttgcgacgaacagcaagttgttctcatcgactgcataaatgtctc
 taacctttccttggcgattaaaggtaggattccgtccagatcagtttcacaatggacatacttgaaggatacagcaaatg
 tgttggagcgtgacacatggaaaggaaattttcgagtttctagtagtagtatattggggcggtgaaagttcagatgttfaatg
 25 ctfaatactcttatactcttcaagcgcccaaggttttctgccaacctgactttttctgaataatgaatcgttcaagtgaggtatta
 aacctgattaaagtacgtgatttggcactggataaggctgaaaaatatccgtattcataaacgattattggtaaaagttaaaa
 ataccactaattacggagaagcttagtaacagttatcatcttggctgaltaacgcttacaatttccattcggcattcaggctgc
 gcaactgttgggaaggcgatcggtgcggccttctcgtattacgccaggcgctcgaggcacaacgaacgtctcactta
 atcttctgtactctgaagaggagtgaggaaataccaagaaaaacatcaaacfcgaatgattttccaaaccctaccacaagat

attcatcagctgcgagataggctgatcaggagcaagctcgtacgagaagaacaaatgacaaaaaaaaatcctatactatat
agggtacaaataaaaaagtatcaaaaatgaagcctgcatctctcaggcgaatggcattctgacatcctcttgattagaatctag
caagaccgggtctctcgtaatgcccactgaactgaggaacagtcagtcctaaaggctacaaactcaatgatgatgatgatg
atggtcgacggcgctattcagatcctctctgagatgagttttgtcggggcccaaaaaataacttcccaatcttcagttatgtcaa
5 gcgafaggttaagaatggatgatcttattatcattctcatatgacgatctagtcggcaatgaactttcttgcatttttttacttta
atactgtcttcaacgitatgttgtgtgtaaatattcttgcgataatgatttttcaatattggtattctcctcaagattgtggatgaat
gcctgtcaaatctcgggaatatgacttgaacatttttaaggtattgtttcgaataaactgaactgagttctactactcacctctt
ggtagccaaaagtctccatcatcaacataaagatctcaacggctcgtccctcgggtatcaggggtataactattacatag
gactgttttcaataacattgaagatctctatacttattctcatagtgataatgtgaccgcttcaataaataatggaattttatcca
10 atgggtgcagaaatattctttctatcaaatcagtagcattgtttataaaagagtgtaaagatgcgaattcatagaatataccgggt
gggaaaaccatttccgttctgattgaccaggctcctgacagggttgactaatagacctgaattactccagtaaaattggttat
cgatatgatacaattcagcaaatcaggggttcaataaacattggattcattatcggaatccagtacacttgatttatgaac
atgggtgataaggtagtagcaggaatatctcaattggataatatcacgtactatcgacttcaagggtcattgaalaagtatg
gttctcttcttgggtgtctataggcggtgtgctctaaaaatgggtaccataagcccgttgggtaccaacgtgcaatcaattca
15 ggtgtaggatcctcagcaagccagctatgctggctcctataaaggcataccagcaatgtgtttgacagaaccataggaat
ggaaatcttaagtaatcccaattggccacattgtcaccagtcctatgtggcagcagtcagtgagagccggcaaaaaagcc
cttgtagaaggaaaggacgttatcggtatggatgaataaatcgattttatgctgctgtaagtagcttcatgactgatagacct
atatgttatggcggatctttctcaatgtaattgtcgtgaatcaaatcttttggagctgtggtctctggccatcgaaaatcgaa
ggctcgttcatatcattccaaatgaataaattagttaaatcagccggcagatccataaaccgttcgaaaaaggacttccaaatc
20 ttttggccataatttgcttatgggtatcaatccatatagaattacctggccagcaatgacctacatagtcatttccattgtgatccttga
ctgctacattttcattaattaccctgtcactgatttcataatcttctttaaattaggatcgattagtagcagaagatttctaccaac
tttttaatttggataacagccttttggattgggaaggagtgctgcttccaagtaaaatatttttgcgttctgtatctccaagtc
caaccaataaaaatcgtaaggaaatcatatgagcatccatctgagagtccactgtgagaacgtccatctcatcattataattccat
ctacattgatgtaccctatagaggaaatggcggtataaaagggtctaccagtcaaatcggtaaatttgcattgatagttgg
25 aatatctgccccaggacatgactacatctatgacaccatttctggagatccaatgagtcacgttttttttactggtgtcata
ctttatgtctaccaagtgtcagctgattgacccaaaagatagatgtggacgaagatgaaacatgaatgggtatgaaccg
tacattgggtgggtggtaccgatgtgtactcaagacatcaacgttgaaaagcctgtaggggtcctttccacctgaagtgtcca
tcagccttagcgacgtcgcattgtccggtataccgtagacattagtagaaccatgaaagagaaatctagcgcaaccgattc
aggcccaaaaggcatagagtcattgttgaatacaagaaattgtccctaaacatgttgaaggtgttcttctggcagcacgtgt

gcgaagtttctgcttagttctatgatgttcaatgttcaggaaatttctcgtttacaataagtttcagcgcatittgccagtaaa
 ctttcaattgaaaaggttcagcaagattttctacggatacatcaccgtttcgaagatgaaatgtgtctgcagtgaggattgaaag
 tgacaaaaatgaagatattttgaccagaatgagttcacagttgttttgcttaaggaagtggaattgtggaactggtcctgt
 tcgctcctcttgaaatttctgtcgaatgcgtacttccagggtcattgaaccgttgaagagatcaacaaccgctgctgttg
 5 gttggcattctctcttctcatttatagtgaaacctactgagtgatcctgtaaaaagagagagagaatgggaactgaacggcta
 tatcatcgccctccaatcttggtatagtttaattaggtagcatgaagcacattcttaaggatcgtgtgcaatagactcggcg
 tccactttgtaatagcagtgatgagatttggaatatttctgcataaacctgtttctatggcaaacccagattgcgcacactt
 ctttaatagatagtcggtaaacgcatgcgaaaaagcggtaagaagaccaattggcatacgagccatttcaaaaggaccatc
 tcg

10

SEQ ID NO: 22 pYPT1ZAGLSII

cgaacaaaaactcatctcagaagaggatctgaatagcggcgtcgaccatcatcatcatcattgagttgtagccttagac
 atgactgttctcagttcaagttgggcacttacgagaagaccggctctgtagattctaataagaggatgacagaatgccattt
 gcctgagagatgcaggctcatitttgatactttttatttgaacctatagtaggattttttgtcattttgttcttctcgtacga
 15 gctgtcctgatcagcctatctcgcagctgatgaatattctgttgtaggggttgggaaatcattcagattgatgttttcttg
 gtatttccactcctcttcagagtacagaagattaagtggacctcgttggcgatccccacacaccatagcttcaaaatg
 ttctactcctttttacttccagatttctcggactccgcgcacgccgtaccacttcaaacacccaagcacagcatactaa
 attttccctcttctcctctagggtgtcgttaattaccgactaaagggttggaaaagaaaaagagaccgcctcgtttcttttc
 ttctgcgaasaaggcaataaaaattttatcacgtttcttttctgaaatttttttttagttttttcttctcagtgacctccattgata
 20 ttaagttaataaacggcttcaatttctcaagttcagtttatttttcttattacaacttttttacttctgttcattagaaagaaa
 gcatagcaatctaataaggggcgggtgttgacaattatcatcggcatagtagtataacgacaaggtgag
 gaaactaaaccatggccaagttgaccagtgcggttccgggtgctaccgcgcgcagctgcgggagcgggtcgagttctgga
 ccgaccgggtcgggttctccgggacttctgtgaggacgacttcggcggtgtggtcgggacgacgtgacctgttcatca
 gcgcgggtccaggaccaggtgtgtcgggacaacacctggcctgggtgtgggtgcgcggcctggacgagctgtacgccg
 25 agtggtcggaggtcgtgtccacgaactccgggacgcctccggcgcccatgaccgagatcggcgagcagccgtggg
 ggcggggagttgccctgcgcgacccggcgcaactgcgtgcacttctgtggcgaggagcaggactgacacgtccgac
 ggcgggccacgggtcccaggcctcggagatccgtcccccttttcttctgcatatcatgtaattagttatgtcacgcttacatt
 cacgccctccccacatccgtctaacccgaaaaggaaggagttagacaacctgaagctaggtccctattttttttatagt
 tatgttagtattaagaacgttatttatatttcaatttttttttctgtacagacgcgtgtacgcatgtaacattatactgaaaacc

ttgcttgagaaggttttgggacgctcgaaggcttaatttgaagctggagaccaacatgtgagcaaaaggccagcaaaag
 gccaggaaccgtaaaaaggccgcgttgcctggcgtttttccataggctcgcggcccccctgacgagcatcacaaaaatcgacgc
 tcaagtcagagggtggcgaacccgacaggactataaagataaccaggcggtttcccccctggaagctccctcgtgcgtcctc
 gttccgaccctgccgttacccggatacctgtccgcctttctcccttcgggaagcggtggcgtttctcaatgctcacgctgtagg
 5 tatlctcagttcgggttaggtcgttcgctccaagctgggcgtgtgcacgaacccccgttcagccccgaccgtgcgccttacc
 cggtaactatcgtcttgagccaacccggtaagacacgacttatcgccactggcagcagccactggtaacaggattagcag
 agcgagggtatgtaggcgggtgctacagagttctgaagtggtggcctaactacggctacacilagaaggacagatttggatct
 gcgctctgctgaagccagttaccttcggaaaaagagttggtagctcttgatccggcaacaaaccaccgctggtagcgggtg
 gttttttgtttgcaagcagcagattacgcgcagaaaaaaggatctcaagaagatcctttgatcttttctacggggtctgacgc
 10 tcagtggaaacgaaaactcacgttaagggaatttgggtcatgagatcagatctatgatgagtcacaatctgcttccacagacgagt
 acaaggacaggcaaaaaggaaattggaagaagtgctaacccaataatgagcaagttctatggagctgctggtggagctcctg
 gtggagctcctggtggcttccctggagggttccctggcggagctggcgcagctggcgggtgcccagggtggtgctgcccga
 ggcgagacagcggaccaaccgtggaagaagtcgattaaagcaattcaacggataaattctggttaatatataacgtgaat
 aggaaattaaggaatttggatctaataatgtgctgtatgccgacatcgggcatcgtagattgtatagatcgtgacactata
 15 ataagccagccaaaaccctaaaccagttgccctccactaattagtgactacccaatcttgctcttcgggtgtctttataag
 gacagattcacaagctcttggcccaatacacacatacacacagagataatagcagtcgaattcacgtggcccagccggc
 cgtctcggatcggtacctcgagatggctcttgaatggctcgtatgccaatggctcttaccgcttttcgcatgcgtttac
 cgactatctattaaagaagtgctcgcaatctgggtttgcatagaacagggttatgcagaaaatattgccaatctcatcac
 tgctattacaagtgagcggagtcattgcacacgatcctttagagaatgtgcttcatgctaccataaataaaactataccaa
 20 gattggaggcgatgatatagccgttcagttccattctctctctttttacaggatcactcagtaagggttcaactataatgag
 aaagagagaatgccaaccaacagcagcggttggatctcttcacaacggttcaatgagacctggaagtacgcattcgaca
 agaaattcaagaggaggcgaacaggaccagttccacaattccacttcttaagcaaaaacaaactgtgaactattctgg
 tcgaaaatatcttattttgtcatttcaaacctccactgcagacacatttcatttcgaacgggtgatgtatccgtagaatctttg
 ctgaacctttcaattgaaagttactggcaaatgcgctgaaactattgtaacgagcaaaatttctgaacattgaacatcat
 25 agaactaagcaggaaaacttcgcacacgtgctgccagaagaacaacttcaacatgttaaggacaatttctgtattcaaa
 gcatgactctatgccttggggcctgaatcgggtgcgctagattctcttcatgggttctactaatgtctacgggtataccggaac
 atgcgacgtcgtgaagctgatggacacttcaggtggaaggaaacctacaggcttttcaacgttgatgtctttgagtacaac
 atcgggtaccagccaaccaatgtacggttcgatccattcatgtttcatcttctccacatctatctttgggtcaatgcagctga
 cacttgggtagacataaagtatgacaccagtaaaaataaacgatgactcattggatctccgaaaatgggtgcatagatgtag

tcatgtccctggggccagatatccaactatcattgacaaattaccgalttgactggtagacccttttaccgcccatttcctcta
 tagggfaccatcaatgtagatggaattataatgatgagatggacgttctcacagtggactctcagatggatgctcatatgattcc
 ttacgattttatttgggtggacttggagtatacgaacgacaaaaaatattttacttgggaagcagcactccttcccaatccaaaaa
 ggctgttatccaaattaaaaagttgggtagaatctgtcgtactaatcgaatcctattaaagaaagattatgaaatcagtgc
 5 agggtaaitaatgaaaatgtagcagtcaggatcacaatggaaatgactatgtaggtcattgctggccaggtaattctatatgg
 attgataccataagcaaatatggccaaaagatttgggaagtccttttcgaacgggttatggatctgccggctgatttaactaattta
 ttcatittggaatgatatgaacgagccttcgattttcgaatggccagagaccacagctccaaaagatttgatcacgacaattac
 attgaggaaagatccgtccataacatatgtgtatcagtgcatgaagctacttacgacgcaataaaatcgatttattcccat
 ccgataagcgtcctttccttcaacaaggccttttttggcggctctcaacgtactgctgccacatggactgggacaatgtggc
 10 caattgggattacttaagatttccattcctatggttctgtcaacaacattgctggatgccatttataggagccgacatagctg
 gcttctgaggatcctacacctgaattgattgcacgttggatccaagcgggcttatggatcccatttttagagcacacgccc
 atalagacaccaagagaagagaaccatactatticaatgaaccttgaagtcgatagtacgtgatattatccaattgagatattt
 cctgctacctaccttataccatgtttcataaatcaagtgtcactggatttccgataatgaatccaatgtttattgaacacctga
 atttgcgaattgtatcatatcgataaccaattttactggagtaattcaggctctattagtcacacctgtcacggagcctggtaac
 15 agaaacggaaatgggtttcccccgggtatattctatgaattcgcactctttactctttataaacaatgggtactgattgataga
 aaagaatatttctgcacattggataaaattccattatttgaaggcgggtcacattatcactatgaaagataagtatagaagat
 cttcaatgttaatgaaaaacgatccatatgtaatagttagccctgataccgaggggacgagccgttgagatctttatgttga
 tgaaggagaaacttttggctaccaaagggtgagtagcgtagaaactcagttcattttcgaaaacaataccttaaaaaatgttcg
 aagtcatattcccagaaatttgacaggcattcaccacaatactttgaggaataccaataattgaaaaaacattatcgaaaagaat
 20 aatttacaacacaataacgttgaagacagtattaaagtcacaaaaaatggcgaagaaagttcattgccgactagatcgtc
 atatgagaatgataataagatcaccattctaacctatcgttgacataactgaagattgggaagttatttttgggcc

SEQ ID NO: 23 pYPT1ADE1glsII

gtcgaccggctgcattaatgaatcggccaacgcgcggggagaggcgggttgcgtattgggcgctcttccgcttcctcgtc
 25 ctgactcgtcgcgtcgtcgttcggctgcggcgagcgggtatcagctcactcaaaggcggtaatacgggttatccacagaatc
 aggggataacgcaggaagaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccgcttgcgtgg
 cgttttccataggtccgccccctgacgagcatcacaaaaatcgacgtcaagtcagagggtggcgaaacccgacagga
 ctataaagataaccaggcgtttccccctggaagctcctcgtgcgtctcctgttccgacctgccgcttaccggatacctgtcc
 gcctttctccctcgggaagcgtggcgtttctcatagctcacgctgtaggtatctcagttcgggtgtaggtcgttcgtccaagc

tgggctgtgtgcacgaacccccgttcagcccgaccgctgcgccttatccggttaactatcgtcttgagtccaacccggttaag
 acacgacttatgccactggcagcagccactggtaacaggattagcagagcgaggatgtagggcgggtctacagagttctt
 gaagtgggtggcctaactacggctacactagaaggacagtatttggtatctgcgctctgctgaagccagttaccttcggaaaaa
 gagttggtagctcttgatccggcaacaaccaccgctggtagcgggtgggttttttgggtgcaagcagcagattacgcgcaga
 5 aaaaaaggatcicaagaagatccttgatctttttacggggtctgacgctcagtggaacgaaaactcacgttaagggattttg
 gtcattgagattatcaaaaaggatctcacctagatccttttaataaaaaatgaagttttaaataatctaaagtatatatgagtaa
 acttgggtctgacagttaccaatgcttaatcagtgaggcacctatctcagcgatctgtctatttcgttcatcatagttgcctgactc
 cccgtcgtgtagataactacgatacgggaggggttaccatctggccccagtgctgcaatgataccgcgagacccacgctca
 ccggctccagattatcagcaataaaccagccagccgggaagggccgagcgcagaagtggctcctgcaactttatccgctcc
 10 atccagctctattaattgttgccgggaagctagagtaagtagtgcgcagttaatagtttgcgcaacgttggtgcatgctacag
 gcatcgtgggtgcacgctcgtcgtttgggtatgggttcattcagctccgggtcccaacgatcaaggcgagttacatgatcccca
 tgttggtcaaaaaagcggttagctccttcggctcctccgatcgttgtcagaagtaagttggccgcagtggtatcactcatggttat
 ggagcagctgcataattcttactgtcatgccatccgtaagatgctttctgtgactgggtgagtactcaaccaagtcattctgag
 aatagtgtatgcggcgaccgagttgctcttgcccggtcaatacgggataataccgcgccacatagcagaactttaaagt
 15 gctcatcattggaaaacgttcttcggggcgaaaactcgaaggatcttaccgctgttgagatccagttcgatgtaaccactcg
 tgcaccaactgatcttcagcatcttttactttaccagcgtttctgggtgagcaaaaacagggaaggcaaaatgccgcaaaaa
 aggggaataaggcgacacgggaatgtgaatactcactcttcttttcaatagctccaaggcaacaattgactactcaga
 ccgacattcattcgttattgattttaataacgataaacggaatgggttacttgaatgatttcatttatgatcattgttactaattac
 ctaaataggattttatatggaattggaagaataagggaatttcagatgtctgaaaaaggcgaggagggtactaatcattcaa
 20 gccatttcttgccagtaattgcttcataagcttcaatafcttttcttactcttgatagcaatttctgcatccatggctacgccctct
 ttgccattcaatccgttggccgtcaaccaatctctgagaaactgcttatcgtactcttgcgatttaccacttggtaagtctttt
 gattccaaaatctagaagaatctggagttaaaactcatctactagtaccaattcattgtttcgtccagtcgaattcgaattcgt
 atcagcaataatgatcccccttcaaaaggcggaagtgttttgagcagaatacaactcgaccgccttgacagcgaccttctcac
 aatgtctttacctacaatctcagcagcttgtcaatagagatgttttcatcgtgttcacctgttcagctttcgttgagggtgtaa
 25 aatcggagttggaaggcgctcgtctctgaagggtctcgttttcaaccttgactccatggacagttttgagttctgtactctttc
 catgcacttccagtgatgtaacctctgacaatggcttccaaaggatcagctctgtgctttttactatcaaggatcgtccctctaat
 tgagatttgtatttttctcagacagttttgatggtagtaagcaagacttcttgcattagaagcaaccaaatgattctttatgta
 ggggtgccaaaaatacaaccagaaaactgagagctgagtcataatcttcccttatcaggaataccgtttgtcataatcacatc
 gtaagcgggagatacggtcagttgcgacgaacagcaagttgtctcatcgactgcataaatgtctctaaccttttcttggcgatt

aaaggtaggattccgtccagatcagtttcacaatggacatacttgaaggatacagcaaagtgtgttgaagcgatgacac
 atggaaaggaattttcagtttcctagagtagtatattggggcgggtgaaagttcagatgtttaatgcitaaactcilitactcttc
 aaagcgcccaagtgtttctgccaacctgactttttctgaataatgaatcgttcaagtggagttttaaacatgattaaagtacgt
 gatttggcactgggataaggtcgaaaaatatccgtattcataaacgattatttggtaaaagttacaaaataccactaattacggag
 5 aagcttagtaacagttatcatctcttggtcgattaacgcttacaattccattcggcattcaggctgcgcaactgttgggaaggg
 cgtatgggtcgggccctcttcgtattacgccaggggcctcgaggcacaacgaacgtctcacttaatctctgtactctgaaga
 ggagtgggaaataccaagaaaaacatcaaacgaatgatttcccaaacccctaccacaagatattcatcagctgcgagat
 aggtgatcaggagcaagctcgtacgagaagaacaaaaatgacaaaaaaatcctatactataggttacaataaaaaa
 gtatcaaaaatgaagcctgcatctctcaggcaaatggcattctgacatcctcttgattagaatctagcaagaccggctctctct
 10 aagtgcccaactgaactgagggaacagtcattgctaaagctacaaactcaatgatgatgatgatgttcgacggcgctatt
 cagatcctctctgagatgagttttgttcgggccccaaaaataacttccaatcttcagttatgtcaagcgataggttaagaatgg
 tgcattatlatcattctcatatgacgatctagtcggcaatgaactttctcgccattttttgactttaactgtcttcaacgttatg
 ttgtgtgtaaaattattcttgcgataatgatttttcaatatttggtattcctcaagattgtggtgaatgcctgtcaaatctcgga
 atatgacttcgaacattttlaaggtattgtttcgaaaatgaactgagtttctacgtactcaccttttggtagccaaaagtttctcc
 15 atcatcaacataaagatctccaacggctcgtccctcggtatcagggggtataactattacatatggatcglttttcattaacattg
 aagatctctatacttattcttcatagtataatgtgaccgccttcaataaataatggaattttatccaatgggtgcagaaatattcttt
 tctatcaaatcagtaccattgtttataaaagagtgaagatgcgaattcatagaatataccgggtgggaaaaccatttccgtttc
 tgattgaccaggetccgtgacaggttgactaatagacctgaattactccagtaaaattggttatcgatatgataaattcagca
 aattcagggtgttcaataaacattggattcattatcggaatccagtgacacttgattatgaaacatgggtataaaggtaggta
 20 gcaggaaatatctcaattggataatcacgtactatcgacttcaaggttcattgaataagtatggttctcttcttgggtctat
 atggggcgtgtgctctaaaaatgggtaccataagcccgttggtaccaacgtgcaatcaattcaggtgtaggatcctcagca
 aagccagctatgtcggctcctataaatggcataccagcaatgttggtagagaaccataggaatggaaatctttaagtaatcc
 caattggccacattgtcaccagtcctatgtggcagcagtagcttgagagccggcaaaaaagcccttgtagaaggaaagga
 cgcttatcggtggtgaataatcgatttattgcgtcgttaagtagcttcatgcactgatagaccatatgttatggacggatctt
 25 tcttcaatgtaattgtcgtgaatcaaatcttttgagctgtggtctctggccatcgaaaatcgaaggctcgttcatatcattcca
 aatgaataaattagttaaatcagccggcagatccataaacggttcgaaaaaggacttccaaatctttggccataatttgcttatg
 gtatcaatccatatagaattacctggccagcaatgacctacatagtcatttccattgtatccttgactgtacatttccattaatta
 cctgtcactgatttcataatcttcttaaatgaggatcatttagcagacaagatttctaccaactttttaattggataacagc
 cttttggattgggaaaggagtgcgtctccaagtaaaatatttttgcgttcgtatactccaagtcacaacaaataaaatcgtaa

ggaatcatatgagcatccatctgagagtcactgtgagaacgtccatctcatcattataattccatctacatigatggfaccctat
agaggaaatggcggtaaaaagggtctaccagtcacaaatcggtaaattgtcaatgafagttggaatatctggccccaggagac
atgactacatctatgacaccaatttcggagatccaatgagtcacgttttttactgggtgcatactttatgtctaccaagtgtc
agctgcattgacccaaaagatagatgtggacgaagatgaaaacatgaatgggatcgaaccgtacattgggtggctggtacc
5 gatgtgtactcaaagacatcaacgtgaaaagcctgtagggttcctttccacctgaagtgtccatcagccttagcgacgtcgc
atgttccggtataccgtagacattagtagaacccatgaaagagaaatctagcgcaaccgattcaggcccaaggcataga
gtcatgcttgaatacaagaaattgtccttaaacatgttgaagttgtttcttctggcagcacgtgtgcgaagtttctgcttagtt
ctatgatgttcaatgttcaggaaatttgcctgtttacaataagtttcacgcattttgccagttaaactttcaattgaaaaggtcag
caaagatttctacggatacatcaccgttcgaagatgaaatgtgtcgcagtgagtttgaagtgcacaaaaatgaagatatit
10 cgaccagaatgagttcacagttgttttgccttaagggaagtgaattgtggaatactggtcctgttcgcctccttgaatttctt
gtcgaatgcgtacttccagggtctcattgaaccgttgtgaagagatcaacaaaccgctgtgttgggtggcattctcttctcat
ttatagtgaaccttactgagtgatcctgtaaaaagagagagagaatgggaactgaacggctatcatcgcctccaatctt
ggtatagttttaattatggtagcatgaagcacattctctaaaggatcgtgtgcaatagactcggcgtccactttgtaafagcagt
gatgagatttggcaatatttctgcataaacctgtttctatggcaaaaccagattgcgcacacttcttaatagatagtcggta
15 aacgcatgcgaaaaagcggtaaagaagaccaattggcatcagagccatttcaaaaggaccatctcagggtaccgatccga
gacggccggtgggcccacgtgaattcgactgctattatctctgtgtgtatgtgtattgggcaacaagagcgttgaatctgt
ccttataaaagacacccgaagaggcaagattgggtagtagactaattagtgaggggcaactgggttaggggtttggctggc
ttattatagtgtcagcgatactatacaatctacgatgccgatgtcggcatacagcacattattagatccaaaatttccttaatttc
ctattcacgttatatataaccagaatttatccgttgaattgcttaatcgacttctccacgggttggtccgctgtctccgctggg
20 gcagcaccacctggggcaccgccagctgcgccagctccgggaaacctccagggaagccaccaggagctccacc
aggagctccaccagcagctccatagaactgtctcattattgggttagcaactcttccaattcctttgcctgtccttgtactcgtc
tgtggaagcagattgtgactcatcatagatctgatctcat

SEQ ID NO: 24 pGAPZAglslIHDEL

25 tcgagatggtccttttgaatggctcgtatgccaattggctcttcttaccgcttttgcgatcgtttaccgactatctattaaagaa
gtgtgcgcaatctgggttttccatagaacagggtttatgcagaaaatttgcacaaatctcatcactgctattacaaagtga
cgccgagctattgcacagatcctttagagaatgtgcttcatgtaccataattaaaactataccaagattggagggcgatga
tatagccgttcagttccattctctctctttttacaggatcactcagtaagggttactataaatgagaaagagagaatgcaa
ccaacagcagcgggttggatctcttcacaacgggttcaatgagacctggaagtacgcattcgacagaagaatticaagagga

ggcgaacaggaccagtattccacaattccacttccttaagcaaaaacaaactgtgaactcattctggtcgaaaatatcttcattt
 ttgtcactttcaaactccactgcagacacatttcatttcgaaacggatgatccgtagaaatcttgcgaacctttcaattg
 aaagtttactggcaaatgcgctgaaacttattgtaaacgagcaaaatttcctgaacattgaacatcatagaactaagcagga
 aaacttcgcacacgtgctgccagaagaacaactttcaacatgtttaaggacaatttcttgattcaaaagcatgactctatgcctt
 5 tggggcctgaatcggtgcgctagatttctcttcatgggttctactaatgtctacgggtataccggaacatgcgacgtcgctaag
 gctgatggacacttcagggtggaagggaacctacaggcttttcaacgttgatgtcttgagtacaacatcggtaccagccaac
 caatgtacgggtcgatcccattcatgtttcatcttcgtccacatctatctttgggtcaatgcagctgacacttgggtagacataa
 agtatgacaccagtaaaaaataaacgatgactcattggatctccgaaatgggtgcatagatgtagtcatgtccctggggcca
 gatattccaactatcatgacaaaattaccgattgactggtagaccttttaccgccatttcctctatagggtaccatcaatgta
 10 gatggaattataatgatgagatggacgttctcacagtggactctcagatggatgctcatatgattccttacgatttatttgggtgg
 acttggagtatacgaacgacaaaaatatttacttgggaagcagcactccttcccaatccaaaagggtgtatccaaattaa
 aaagtgggtagaatctgtcgtactaatcgtactcatttaaagaaagattatgaaatcagtgacagggtaatatgaaat
 gtagcagtcaaggatcacaatggaaatgactatgtaggtcattgctggccagggttaattctatgattgataaccataagcaaa
 tatggccaaaagatttgggaagtccttttcgaacgggttatggatctgccggctgatttaactaatttattcatttgggaatgatatga
 15 acgagccttcgatlttcgatggcccagagaccagctccaaaagatttgattcacgacaattacattgaggaaagatccgtc
 cataacatataatggctatcagtgcatgaagctactacgacgcaataaaatcgaatttaccatccgataagcgtccttctc
 tctaacaagggttttttgcggctctcaacgtactgctgccacatggactgggtgacaatgtggccaattgggattacttaag
 atttccattcctatgggtctgtcaacaacattgctggatgccatttataggagccgacatagctggcttctgaggatcctac
 acctgaattgattgcacgttggtagcaagcgggcttatggtagccatttttagagcacacgcccatatagacaccaagagaa
 20 gagaaccatactattcaatgaacctttgaagtcgatagtacgtgatattatccaattgagatatttctgctacctaccttataca
 ccatgttcataaatcaagtgactggttccgataatgaatccaatgtttattgaacacctgaatttctgaattgtatcatat
 cgataaccaattttactggagtaattcagggtctattagtaaacctgtcacggagcctgtcaatcagaacggaaatggtttc
 ccacccgggtatattctatgaattcgcactctttataacaatggtagtattgatagaaaagaatttctgcaccat
 tggataaaattccattatttattgaaggcgggtcacattatcactatgaaagataagtatagaagatcttcaatgttaataaaaaac
 25 gatccatatgtaatagttagccctgataccgaggagcagccgttggagatctttatgttgatgatggagaacttttggct
 accaaagagggtgagtacgtagaactcagttcattttgaaaacaataccttaaaaaatgttcgaagtcatttcccgagaattt
 gacaggcatccaccacaatactttgaggaataccaatattgaaaaatcattatcgcaaagaataatttacaacacaacataac
 gttgaaagacagtattaaagtaaaaaaatggcgaagaaagttcattgccgactagatcgtcatatgagaatgataataaga
 tcaccattcttaacctatcgcttgacataactgaagattgggaagtttttgggccgaacaaaaactcatctcagaagagg

atctgaatagcgcgcgtgaccacgacgaactgtgagtttttagcccttagacatgactgttccctcagttcaagttggggcacttacg
agaagaccgggtcttgtagatttctaatacaagggaatgcagaatgccattgacctgagagatgcaggcttcatttttgatactttt
ttatttgaacctataatagtataggattttttgtcattttgttctctctgacgagcftgctcctgatcagcctatctcgcagctgat
gaataatcttggtgaggggtttgggaaatcattcagatttgatgtttttcttggtatttccactcctcttcagagtacagaagatt
5 aagtgagaccttcgtttgfcggatccccacacaccatagcttcaaaatgtttctactcctttttactcttccagattttctcgga
ctccgcgcacgcgtaccacttcaaaacaccaagcacagcatactaaattttccctcttttcttctctaggggtgctgtaatta
cccgtactaaagtttggaaaagaaaaagagaccgcctcgtttttttcttctgctgaaaaaggcaataaaaaattttatcacg
tttctttttctgaaaatttttttttagttttttcttcttccagtacccctcattgataatttaagtaataaacggcttcaatttctcaagttt
agtttcatittttctgttctattacaacttttttacttctgttctcattagaaagaaagcatagcaatctaataaggggcggtgttgac
10 aattaatcatcggcatagtatcggcatagataatacagacaagggtgaggaaactaaaccatggccaagttgaccagtgccgt
tccggtgctcaccgcgcgcgacgtcgcggagcggctcagttctggaccgaccggctcgggttctccgggacttcgtgg
aggacgacttcgccggtgtggtccgggacgacgtgacctgttcatcagcgcggtccaggaccaggtggtgccggacaa
caccctggccctgggtgtgggtgcgcggcctggacgagctgtacgccgagtggtcggaggtcgtgtccacgaactccgg
gacgcctccggggccggccatgaccgagatcggcgagcagccgtgggggagggttcgcccgtgcgcgacccggccg
15 gcaactgcgtgcacttcgtggccgaggagcaggactgacacgtccgacggcgccacgggtcccaggcctcggagat
ccgtcccccttttcttctgtcgtatcatgtaattagttatgtcacgcttaccacgcccctccccccacatccgctctaaccgaa
aagggaaggagttagacaacctgaagtctaggtccctattttttttatagttatgttagttaagaacgtatttatatttcaaatt
tttcttttttctgtacagacgcgtgtacgcatgttaacttatactgaaaaccttgcttgagaaggttttgggacgctcgaaggct
ttaatttgcaagctggagaccaacatgtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccgcttgcgtgg
20 cgtttttccataggtccgccccctgacgagcatcacaaaaatcgacgctcaagtcagaggtggcgaaacccgacagga
ctataaagataccaggcgtttccccctggaagctccctcgtgcgtctcctgttccgacctgcccgttaccggatacctgtcc
gcctttctcccttcgggaagcgtggcgcttttcaatgtctacgctgtaggtatctcagttcgggtgtaggtcgttctccaagc
tgggctgtgtgcagcaacccccgttcagcccagccgctgcgccttatccggtaactatcgtcttgagtccaacccggtaag
acacgacitattgccactggcagcagccactggttaacaggattagcagagcgagggtatgtaggcgggtgtacagagttctt
25 gaagtggtagctcttgatccggcaaacaaaccaccgctggtagcgggtgttttttgtttgcaagcagcagattacgcgcaga
aaaaaaggatctcaagaagatcctttgatcttttctacggggtctgacgctcagtggaacgaaaactcacgttaagggaatttg
gtcatgcatgagatcagatcttttttagaaatgtcttgggtgtcctcgtccaatcaggtagccatcttgaatatctggctccgt
tgcaactccgaacgacctgctggcaactgaaaattctccggggtaaaaacttaaatgtggagtaattggaaccagaaacctctc

ttccctctctctcctccaccgccggtaccgtccctaggaaatttactctgctggagagcttctctacggcccccttgacgc
aatgctcttcccagcattacgttgcgggtaaaacggaggctgtgtaccggacctagcagcccagggtggaaaagtcccg
gccgtcgtggcaataatagcggggcgacgcattgcatgagatttggaaaccaccagaatcgaatataaaaggcgaac
accttcccaatttgggttctcctgacccaagactttaatttaatttatttgcctatttcaatcaattgaacaactatttcgaaa
5 cgaggaattcacgtggcccagccggcgtctcggtacc

SEQ ID NO: 25 pGAPADE1glsIIHDEL

cgctactcacctcttggtagccaaaagtctccatcatcaacataaagatctccaacggctcgtccctcggtatcaggggctat
aactattacataggtacgttttccattgaagatcttatacttcttcatagtgataatgtaccgacctcaataaataa
10 tggaaatttccaatggtgcagaaatattctttctatcaaatcagttacattgtttataaaagagtgtaaagatcggaattcatag
aatataccgggtgggaaaaaccatttccgttctgattgaccaggctccgtgacagggttgactaafagacctgaattactccag
taaaattgggtatcgatatgatacaattcagcaaatcagggtgttcaataaacattggattcattatcggaatccagtacact
tgatttatgaaacatggtgtataaggtaggtagcaggaaatctcaattggataaatcacgtactatcgacticaaagggtcat
tgaataagtatggttctcttcttgggtctatatgggcgtgtgctctaaaaatgggtaccataagcccgttggtaccaacgt
15 gcaatcaattcaggtgtaggatcctcagcaagccagctatgtcggctcctataaatggcataccagcaatgttgttgacag
aaccataggaatggaaatcttaagtaatcccaattggccacattgtcaccagtcctgtggcagcagtcagttgagagccg
gcaaaaaagccctgttagaaggaaaggacgcttatcggtggtgaataaacgattttatgcgtcgtagtagcttcatgc
actgatagaccatatgttatggacggatcttctcaatgtaattgtcgtgaatcaaatctttggagctgtggtctctgggcca
tcgaaaatcgaaggctggtcatatcattccaaatgaataaattagttaaatcagccggcagatccataaaccgttcgaaaaag
20 gacttccaaatctttggccatatttgcctatggtatcaatccatatagaattacctggccagcaatgacctacatagtcatttccat
tgtgatccttgactgtacattttcatttaattaccctgtcactgatttcataatcttctttaaagaggatcgatttagtacgacaaga
tttctaccaacttttttaatttggataacagccttttggattgggaaaggagtgtgcttccaagtaaaatattttgtcgttcgta
tactccaagtccaaccaataaaatcgtgaaggaatcatatgagcatccatctgagagtcactgtgagaacgtccatctcatc
attataattccatctacatigatgggtaccctatagaggaaatggcggtaaaaagggtctaccagtcaaatcggtaaatttgc
25 atgatagttggaaatctggccccaggacatgactacatctatgacaccatttccggagatccaatgagtaicgtttattttta
ctggtgtcactttatgtctaccaagtgtcagctgcattgacccaaaagatagatgtggacgaagatgaaaacatgaatgg
gatcgaaccgtacattgggtggctgtgaccgatgttactcaagacatcaacgttgaaaagcctgtagggttcccttccacc
tgaagtgtccatcagccttagcgacgtcgcatgttccggtataccgtagacattagtagaaccatgaaagagaaatctagc
gcaaccgattcagggcccaaggcatagatgcttgaatacaagaaattgtccttaaacatgttgaaggtgtttctctg

gcagcacgtgtcgaagtttccctgcttagttctatgatgttcaatgttcaggaaatttgcctggttacaataagtttcagcgcatt
 ttgccagtaaacatttcaattgaaaagggttcagcaaagatttctacggatacatcaccgtttcgaagatgaaatgtgtctgcagtg
 gagtttgaaagtgcacaaaaatgaagatatttgcaccagaatgagttcacagtttgttttgcittaagggaagtggattgtggaat
 actggctcctgttcgcctcctcttgaatttctgtcgaatgcgtacttccaggtctcattgaaccgttgaagagatcaacaac
 5 cgctgctgttggttggcatttctcttctcatttatagtgaaccttactgagtgatcctgtaaaaaagagagagagaatgggaa
 ctgaacggctatatcatcgccctccaatcttggtatagtttaattatggtagcatgaagcacattctctaaaggatcgtgtgcaa
 tagactcggcgtccactttgtaatagcagtgatgagatttggcaatatttctgcataaacctgtttctatggcaaaaccagat
 tgcgcacacttcttaatagatagtcggtaaacgcgtgcgaaaaagcggtaagaagaccaattggcatacgagccatttca
 aaaggaccatctcgaggtaccgatccgagacggcggctggccacgtgaattcctcgttgcgaaatagttgtcaattgatt
 10 gaaataggagacaataaataaattaaagtcttgggtcaggagaaacaaaattgggaaagggtgttcgcttttatattcgtat
 tcttggtgtgttccaataatctcatgacatgcgtccgccgtatttattgccagcgacggccgggacttttccatccctgggctg
 cttaggtcgggtacacgacctccgtttaccgcgaacgtaatgctgggaagagcattgctgcaagggggccgtagaagaag
 ctctccagcagagtaaaatttcttagggacggtaacggcggtggaaggagagagaagggaagagacgttctgttcca
 ttactccacatttaagtttaccgggagaatttaccgttgcagcaggtcgttcggagttgcaacggagccagatatttcagag
 15 atggctacctgattggacgaggacaccaagacatttctacaaaaagatctgatctcatcgaccggctgcattaatgaatcgg
 ccaacgcgcggggagagggcgggttgcgtattggcgctcttccgttccctcgtcactgactcgtcgcctcggctgttcgg
 ctgcggcgagcgggtatcagctcactcaaggcggtaatacggttatccacagaatcaggggataacgcaggaagaacat
 gtgagcaaaaggccagcaaaaggccaggaaccgtaaaaaggccgctgtgctggcgttttccataggtccgccccctg
 acgagcatcacaaaaatcgacgtcaagtcagaggtggcgaaaccgcagagactataaagataccaggcgtttccccct
 20 ggaagctccctcgtgcgtctcctgttccgaccttccggttaccggatacctgtccgcttctccttccgggaagcgtggc
 gctttctcatagctcacgctgtaggtatctcagttcgggtgtaggtcgttcgctccaagctgggctgtgtgcacgaacccccgt
 tcagcccgaccgtcgccttatccgtaactatcgtcttgagtccaaccggtaagacacgacttatcgccactggcgaca
 gccacttggtaacaggattagcagagcgaggtatgtaggcgggtgtacagagttcttgaagtgggtgcctaactacggctac
 actagaaggacagtatttggatctgcgtctgctgaagccagttaccttcggaaaaagagttggtagctcttgatccggcaa
 25 acaaacaccgctggtagcgggtgtgtttttgttgaagcagcagattacgcgcagaaaaaaggatctcaagaagatcctt
 tgatcttttctacggggtctgacgctcagtggaacgaaaactcacgttaagggaatttggcatgagattatcaaaaaggatctt
 cacctagatccttttaattaaaaatgaagtitttaaatcaatctaaagtatatatgagtaaaacttggctgacagttaccaatgctta
 atcagtgaggcacctatctcagcgtatctgtatttctgtatccatagttgcctgactccccgtcgttagataactacgatac
 gggagggccttaccatctggccccagtgctgcaatgataccgcgagaccacgctcaccggctccagatttatcagcaataa

accagccagccggaaggccgagcgcagaaagtggtcctgcaactttatccgcctccatccagctctattaattgttgcggg
aagctagagtaagtagttccagttaatagtttgcgcaacgttggtgccattgctacaggcatcgtggtgtcacgctcgtcgtt
tggtatggcttcattcagctccggtcccaacgatcaaggcgcagttacatgatccccatggtgtgcaaaaaagcggtagctc
cttcggtcctccgatcgtgtcagaagtaagttggccgcagtggtatcactcatggttatggcagcactgcataattctctactg
5 tcatgccatccgtaagatgcttttctgtgactggtagtactcaaccaagtcattctgagaatagtgtatgcggcgaccgagtt
gctcttccccggcgtcaatacgggataataccgcgccacatagcagaactttaaagtgctcatcattggaaaacgttctcg
ggcgaaaaactctcaaggatcttaccgctgttgagatccagttcgatgaaccactcgtgcaccaactgatcttcagcatct
tttactttcaccagcgtttctgggtgagcaaaaacaggaaggcaaaatgcgcgcaaaaagggaataaggcgacacggaa
atgttgaatactcatactcttcttttcaatagctccaaggcaacaaatgactactcagaccgacattcattcgttatigatttaa
10 atcaacgataaacggaatggttacttgaatgatttactttatgatcattgtttactaattacctaataatagagttttatatggaattgg
aagaataagggaatttcagatgtctgaaaaaggcgaggagggtactaatcattcaagccatttcttgcagtaattgcttca
taagcttcaatafacttttcttactcttgatagcaatttctgcatccatggctacgccctcttggcattcaatccgttggcgtcaa
ccaatctctgagaaactgcttactgtaactctcttgcgatttaccacttggtaagtcttttgattccaaaatctagaagaatctgg
agttaaaactcatctactagtaccaattcattgtttcgtccagttcaaattcgaatttctatcagcaataatgatccccitcaaa
15 agggcgaaagtttttgcagcagaataacaactcgaccgcttgacagcgaccttctcacaatgtctttacctacaatctcagca
gcttgttcaatagagatgtttcatcgtgttcaccctgttcagcttctgtgaagggtgtaaaatcgaggttgaaaggcgtcgtc
ctcttgaagggtctcgtttcaaccttgactccatggacagtgtttgagttctgtactcttccatgcacttccagtgtgtaacctc
tgacaatggcttccaaaggatcagctctgtctttttactatcaaggatcgtccctctaattgagatttgtttttctcagacagt
tttgatggtagtaaagcaaaagacttctgtcattagaagcaaccaaattgattctttatgtagggtgcaaaaaatcaaacag
20 aaaactgagagctgagtcataatcttcccttatcaggaataccggttgcataatcacatcgtgaagcggagatacggtcagtt
gcgacgaacagcaagttgttctcatcgactgcataaatgtcttaaccttcttggcgattaaaggtaggattccgtccagat
cagtggtcacaatggacatacttgaaggatacagcaagtggttgaagcgtgacacatggaaaggaaattttcagttt
cctagagtagtatattggggcgggtgaaagttcagatgtttaatgcttaatactcttatacttcaagcgcccaaggtttctgc
caacctgactttttctgaataatgaatcgttcaagtgagattttaaaccatgattaagttacgtgatttggcactggataaggct
25 gaaaaatatccgtatcataaacgattattggtaaaagttacaaaataccactaattacggagaagcttagtaacagttatcatct
cttggctgattaacgcttacaatttccattcgccattcaggctgcgcaactgttgggaaggcgatcgggtcgggcctcttcg
ctattacgccaggcgctcgaggcacaacgaacgtctcacttaattcttctgactctgaagaggagtgaggaaataccaagaa
aaacatcaaaactcgaatgattttccaaacccctaccacaagatattcatcagctgcgagataggctgatcaggagcaagct
cgtacgagaagaacaaaatgacaaaaaaatcctafactataggttacaaaataaaaaagtatcaaaaatgaagcctgca

tctctcaggcaaatggcattctgacatcctcttgattagaatctagcaagaccggctctctcgttaagtgcccaactgaactgag
gaacagtcattgtctaaggctaaaactcacagttcgtcgtggcgacggcgctattcagatcctctctgagatgagttttgttc
gggccccaaaaataactcccaatcttcagttatgtcaagcgatagggttaagaatgggatcttattatcattctcatatgacgatc
tagtcggcaatgaactttctcgccattttttgactttaatactgtcttcaacgttatgttggttgtaaattattcttgcgataatg
5 atttttcaatattgggtattcccaagtattgtgggaatgcctgtcaaatctcgggaatatgacttcgaacatttttaaggtattg
ttttcgaaaatgaactgagtttcta

SEQ ID NO: 26

GACGAGATCTTTTTTTCAGACCATATGACCGG

10

SEQ ID NO: 27

GCGGAATTCTTTTCTCAGTTGATTTGTTTGT

SEQ ID NO: 28

15 GCG GGT CGA CCA CGA CGA ACT GTG AGT TTT AGC CTT AGA CAT GAC

SEQ ID NO: 29

CAG GAG CAAA GCT CGT ACG AG

20

We claim:

1. A vector capable of expressing an α -1,2-mannosidase or a functional part thereof in a methylotrophic yeast strain, comprising a nucleotide
5 sequence coding for said α -1,2-mannosidase or said functional part.
2. The vector of claim 1, wherein said α -1,2-mannosidase is a protein from a fungal species.
- 10 3. The vector of claim 2, wherein said fungus is *Trichoderma reesei*.
4. The vector of claim 1, wherein said α -1,2-mannosidase is a protein from a mammalian species.
- 15 5. The vector of claim 4, wherein said α -1,2-mannosidase is murine α -1,2-mannosidase IA or IB.
6. The vector of claim 1, wherein said α -1,2-mannosidase or said functional part is tagged with an ER-retention signal.
20
7. The vector of claim 6, wherein said ER-retention signal comprises peptide HDEL.
8. The vector of claim 1, wherein the nucleotide sequence coding for
25 said α -1,2-mannosidase or said functional part is operably linked to a promoter and a 3' termination sequence.

9. The vector of claim 8, wherein said promoter is the promoter of a gene selected from the group consisting of AOXI, AOXII, GAP, and FLD.

10. A vector selected from the group consisting of
5 pGAPZMFManHDEL, pGAPZMFManMycHDEL, pPICZBMFManMycHDEL, pGAPZmManHDEL, pGAPZmMycManHDEL, pPIC9mMycManHDEL and pGAPZmMycManHDEL.

11. A vector capable of expressing a glucosidase II or a functional part
10 thereof in a methylotrophic yeast strain, comprising a nucleotide sequence coding for said glucosidase II or said functional part.

12. The vector of claim 11, wherein said glucosidase II is a protein
from a fungal species.

15

13. The vector of claim 12, wherein said fungus is *Saccharomyces cerevisiae*.

14. The vector of claim 11, wherein said glucosidase II is a protein
20 from a mammalian species.

15. The vector of claim 11, wherein said glucosidase II or said functional part is tagged with an ER-retention signal.

25 16. The vector of claim 15, wherein said ER-retention signal comprises peptide HDEL.

17. The vector of claim 11, wherein the nucleotide sequence coding for said α -1,2-mannosidase or said functional part is operably linked to a promoter and a 3' termination sequence.
- 5 18. The vector of claim 17, wherein said promoter is the promoter of a gene selected from the group consisting of AOXI, AOXII, GAP, and FLD.
19. A vector having the designation pGAPZAGLSII, pPICZAGLSII, pAOX2ZAGLSII, pYPTIZAGLSII, pGAPADEglsII, pPICADEglsII,
10 pAOX2ADEglsII, pYPTIADEglsII, pGAPZAglsIIHDEL and pGAPADEglsIIHDEL.
20. A vector for disrupting the Och1 gene in a methylotrophic yeast strain, comprising a portion of the Och1 gene and a selectable marker gene, wherein said portion of the Och1 gene and said selectable marker gene are linked in such a way
15 to effect the disruption of the genomic Och1 gene in said methylotrophic yeast strain.
21. A vector having the designation pBLURA5'PpOCH1.
22. A method of reducing the glycosylation on proteins produced from
20 a methylotrophic yeast, comprising transforming said yeast with any one of the vectors of claims 1-21.
23. The method of claim 22, wherein said yeast is *Pichia pastoris*.
- 25 24. The method of claim 23, wherein said yeast is a *Pichia pastoris* strain selected from GS115 (NRRL Y-15851), GS190 (NRRL Y-18014), PPF1 (NRRL Y-18017), PPY12-OH, yGC4, or derivatives thereof.

25. The method of claim 22, 23 or 24, wherein said yeast has been genetically engineered to expresses a heterologous protein.

26. A genetically engineered strain of a methylotrophic yeast, wherein
5 said strain is transformed with at least one of the vectors of claims 1-21.

27. A method of reducing the glycosylation of a heterologous glycoprotein expressed from a methylotrophic yeast, comprising transforming cells of said methylotrophic yeast with at least one of the vectors of claims 1-21, and
10 producing said glycoprotein from the transformed cells.

28. A method of producing a glycoprotein with reduced glycosylation in a methylotrophic yeast, comprising transforming cells of said methylotrophic yeast with at least one of the vectors of claims 1-21 and with a nucleotide sequence capable
15 of expressing said glycoprotein in said yeast, and producing said glycoprotein from the transformed cells.

29. A glycoprotein produced by the method of claim 27 or 28.

20 30. The glycoprotein of claim 29, wherein said glycoprotein has a reduced immunogenicity as relative to the glycoprotein produced from a wild type strain of said methylotrophic yeast.

31. The glycoprotein of claim 29, wherein said glycoprotein is suitable
25 for use in human therapeutics.

32. A kit comprising any of the vectors of claims 1-21.

33. The kit of claim 32, further comprising a methylotrophic yeast strain.

34. A kit comprising the methylotrophic yeast strain of claim 26.

5




FIGURE 1

L929 (cycloheximide+ Ti^{-} induced) mouse cDNA library,
Group 6, average insert length : 2000 bp.




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8

Pichia OCH1 Genomic fragment




FIGURE 9




FIGURE 10




FIGURE 11

GlsII Pichia expression vector pGAPZ(A)glsII




FIGURE 12




FIGURE 13

GlsII Pichia expression vector pPICZ(A)glsII




FIGURE 14

GlsII Pichia expression vector pYPT1Z(A)glsII




FIGURE 15

GlsII Pichia expression vector pGAPADE1glsII




FIGURE 16

GlsII Pichia expression vector pAOX2ADE1glsII




FIGURE 17

GlsII Pichia expression vector pPICADE1glsII




FIGURE 18

GlsII Pichia expression vector pYPT1ADE1glsII




FIGURE 19

GlsII Pichia expression vector pGAPZ(A)glsIIHDEL




FIGURE 20

GlsII Pichia expression vector pGAPADE1glsIIHDEL




FIGURE 21

Glucosidase II assay on commercially available alpha-glucosidase




FIGURE 22

Glucosidase II assay on heterologously expressed Pichia protein



FIGURE 23

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2002 (03.01.2002)

PCT


(10) International Publication Number
WO 02/000856 A3

- (51) International Patent Classification⁷: C12N 15/80, 15/62, 15/56, 9/24, C12P 21/00
- (21) International Application Number: PCT/IB01/01514
- (22) International Filing Date: 29 June 2001 (29.06.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/215,676 30 June 2000 (30.06.2000) US
- (71) Applicant: FLANDERS INTERUNIVERSITY INSTITUTE FOR BIOTECHNOLOGY (VIB) [BE/BE]; Rijvisschestraat 120, B-9052 Zwijnaarde (BE).
- (72) Inventors: CONTRERAS, Roland; Molenstraat 53, B-9820 Merelbeke (BE). CALLEWAERT, Nico, L.,
- M.; Ketelbuiserstraat 99, B-8810 Lichtervelde (BE). GEYSENS, Steven, C., J.; Steenbeekstraaje 8, B-9770 Kruishoutem (BE).
- (74) Agent: PUST, Detlev, A.; Hoffmann . Bitle, Arabellas-trasse 4, 81925 München (DE).
- (81) Designated States (*national*): AU, CA, JP, KR, MX, NO.
- (84) Designated States (*regional*): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).
- Published:
— with international search report
- (88) Date of publication of the international search report:
3 January 2003

[Continued on next page]

(54) Title: PROTEIN GLYCOSYLATION MODIFICATION IN *PICHA PASTORIS*

GlsII *Pichia* expression vector pGAPADE1glsIIHDEL



(57) Abstract: The present invention provides genetically engineered strains of *Pichia* capable of producing proteins with reduced glycosylation. In particular, the genetically engineered strains of the present invention are capable of expressing either or both of an α -1,2-mannosidase and glucosidase II. The genetically engineered strains of the present invention can be further modified such that the OCH1 gene is disrupted. Methods of producing glycoproteins with reduced glycosylation using such genetically engineered strains of *Pichia* are also provided.

WO 02/000856 A3



For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

International Application No.

CT/IB 01/01514

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/80 C12N15/62 C12N15/56 C12N9/24 C12P21/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C12P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

WPI Data, PAJ, CAB Data, SEQUENCE SEARCH, BIOSIS, EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category ^a

Citation of document, with indication, where appropriate, of the relevant passages

Relevant to claim No.

X

MARTINET W ET AL: "MODIFICATION OF THE
PROTEIN GLYCOSYLATION PATHWAY IN THE
METHYLOTROPHIC YEAST PICHIA PASTORIS"
BIOTECHNOLOGY LETTERS, KEW, SURREY, GB,
vol. 20, no. 12, December 1998 (1998-12),
pages 1171-1177, XP001006335
ISSN: 0141-5492

1-3, 6, 8,
9, 29-31

Y

cited in the application
the whole document

6.7

Y

EP 0 548 012 A (CIBA GEIGY AG)
23 June 1993 (1993-06-23)
the whole document

6.7

— / —

☒

Further documents are listed in the continuation of box C.

☒

Patent family members are listed in annex.

° Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

17 September 2002

Date of mailing of the International search report

08. 10. 2007

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer _____

Hornig, H

INTERNATIONAL SEARCH REPORT

National Application No.

PCT/IB 01/01514

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	NAKAYAMA K-I ET AL: "OCHI ENCODES A NOVEL MEMBRANE BOUND MANNOSYLTRANSFERASE: OUTER CHAIN ELONGATION OF ASPARAGINE-LINKED OLIGOSACCHARIDES" EMBO JOURNAL, OXFORD UNIVERSITY PRESS, SURREY, GB, vol. 11, no. 7, 1992, pages 2511-2519, XP002934332 ISSN: 0261-4189 figure 3	20
E	EP 1 211 310 A (KIRIN BREWERY ;NAT INST OF ADVANCED IND SCIEN (JP)) 5 June 2002 (2002-06-05) page 14, line 46 - line 56	20
X	KNISKERN P J ET AL: "CHARACTERIZATION AND EVALUATION OF A RECOMBINANT HEPATITIS B VACCINE EXPRESSED IN YEAST DEFECTIVE FOR N-LINKED HYPERGLYCOSYLATION" VACCINE, BUTTERWORTH SCIENTIFIC. GUILDFORD, GB, vol. 11, no. 12, 1994, pages 1021-1025, XP001073955 ISSN: 0264-410X the whole document	29-31
Y	TROMBETTA E SERGIO ET AL: "Endoplasmic reticulum glucosidase II is composed of a catalytic subunit, conserved from yeast to mammals, and a tightly bound noncatalytic HDEL-containing subunit." JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 271, no. 44, 1996, pages 27509-27516, XP002213792 ISSN: 0021-9258 the whole document	11-18
Y	CREGG J M ET AL: "High-level expression and efficient assembly of Hepatitis B surface antigen in the methylotrophic yeast, Pichia pastoris" BIO/TECHNOLOGY, NATURE PUBLISHING CO. NEW YORK, US, vol. 5, 1 May 1987 (1987-05-01), pages 479-485, XP002087851 ISSN: 0733-222X the whole document	11-18
Y	WO 91 05057 A (SALK INST BIOTECH IND) 18 April 1991 (1991-04-18) the whole document	11-18
	-/-	

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/IB 01/01514

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	NAKANISHI-SHINDO Y ET AL: "STRUCTURE OF THE N-LINKED OLIGOSACCHARIDES THAT SHOW THE COMPLETE LOSS OF ALPHA-1,6-POLYMANNOSE OUTER CHAIN FROM OCH1, OCH1 MNN1, AND OCH1 MNN1 ALG3 MUTANTS OF SACCHAROMYCES CEREVISIAE" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 268, no. 35, 15 December 1993 (1993-12-15), pages 26338-26345, XP002934334 ISSN: 0021-9258 the whole document	
A	LEHLE L ET AL: "GLYCOPROTEIN BIOSYNTHESIS IN SACCHAROMYCES CEREVISIAE: NGD29, AN N-GLYCOSYLATION MUTANT ALLELIC TO OCH1 HAVING A DEFECT IN THE INITIATION OF OUTER CHAIN FORMATION" FEBS LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 370, no. 1/2, 1995, pages 41-45, XP001018406 ISSN: 0014-5793 the whole document	
A	EP 0 582 244 A (BOEHRINGER MANNHEIM GMBH) 9 February 1994 (1994-02-09) the whole document	
A	YOKO-O A T ET AL: "Schizosaccharomyces pombe och1 encodes alpha-1,6-mannosyltransferase that is involved in outer chain elongation of N-linked oligosaccharides" FEBS LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 489, no. 1, 26 January 2001 (2001-01-26), pages 75-80, XP004239401 ISSN: 0014-5793 the whole document	
A	LAI A ET AL: "SUBSTRATE SPECIFICITIES OF RECOMBINANT MURINE GOLGI ALPHA 1,2-MANNOSIDASES IA AND IB AND COMPARISON WITH ENDOPLASMIC RETICULUM AND GOLGI PROCESSING ALPHA1,2-MANNOSIDASES" GLYCOBIOLOGY, IRL PRESS, GB, vol. 8, no. 10, October 1998 (1998-10), pages 981-995, XP000952920 ISSN: 0959-6658 the whole document	

-/--

INTERNATIONAL SEARCH REPORT

national Application No

PCT/IB 01/01514

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	MARAS MARLEEN ET AL: "Molecular cloning and enzymatic characterization of a <i>Trichoderma reesei</i> 1,2-alpha-D-mannosidase." JOURNAL OF BIOTECHNOLOGY, vol. 77, no. 2-3, 17 February 2000 (2000-02-17), pages 255-263, XP002205095 ISSN: 0168-1656 cited in the application the whole document ---	
A	HERSCOVICS A ET AL: "ISOLATION OF A MOUSE GOLGI MANNOSIDASE CDNA, A MEMBER OF A GENE FAMILY CONSERVED FROM YEAST TO MAMMALS" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 269, no. 13, 1 April 1994 (1994-04-01), pages 9864-9871, XP002946047 ISSN: 0021-9258 cited in the application the whole document ---	
A	LAL A ET AL: "ISOLATION AND EXPRESSION OF MURINE AND RABBIT CDNAS ENCODING AN ALPHA1,2-MANNOSIDASE INVOLVED IN THE PROCESSING OF ASPARAGINE-LINKED OLIGOSACCHARIDES" JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 269, no. 13, 1 April 1994 (1994-04-01), pages 9872-9881, XP002946050 ISSN: 0021-9258 cited in the application the whole document ---	
A	TREMBLAY L O ET AL: "CLONING AND EXPRESSION OF A SPECIFIC HUMAN ALPHA,2-MANNOSIDASE THAT TRIMS MAN9GLCNAC2 TO MAN8GLCNAC2 ISOMER B DURING N-GLYCAN BIOSYNTHESIS" GLYCOBIOLOGY, IRL PRESS,, GB, vol. 9, no. 10, October 1999 (1999-10), pages 1073-1078, XP000952919 ISSN: 0959-6658 the whole document ---	

-/-

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 01/01514

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>GONZALEZ DANIEL S ET AL: "Identification, expression, and characterization of a cDNA encoding human endoplasmic reticulum mannosidase I, the enzyme that catalyzes the first mannose trimming step in mammalian Asn-linked oligosaccharide biosynthesis"</p> <p>JOURNAL OF BIOLOGICAL CHEMISTRY, AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 274, no. 30, 23 July 1999 (1999-07-23), pages 21375-21386, XP002150706 ISSN: 0021-9258 the whole document</p> <p>----</p>	
A	<p>WO 96 21038 A (MARAS MARLEEN ; CONTRERAS ROLAND (BE); ALKO GROUP LTD (FI)) 11 July 1996 (1996-07-11) the whole document</p> <p>----</p>	
A	<p>EP 0 314 096 A (ZYMOGENETICS INC) 3 May 1989 (1989-05-03) the whole document</p> <p>----</p>	
T	<p>CALLEWAERT NICO ET AL: "Use of HDEL-tagged Trichoderma reesei mannosyl oligosaccharide 1,2-alpha-D-mannosidase for N-glycan engineering in Pichia pastoris."</p> <p>FEBS LETTERS, vol. 503, no. 2-3, 17 August 2001 (2001-08-17), pages 173-178, XP002205096 ISSN: 0014-5793 the whole document</p> <p>-----</p>	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/IB 01/01514

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claim 17 was read as:

17. The vector of claim 11, wherein the nucleotide sequence coding for said glucosidase II or said functional part is operably linked to a promoter and a 3' termination sequence.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: (1-10)-complete, (22-24)-partially

A vector capable of expressing an alpha-1,2-mannosidase or a functional part thereof in a methylotrophic yeast strain, comprising a nucleotide sequence coding for said alpha-1,2-mannosidase or said functional part; a vector selected from the group consisting of pGAPZMFMHDEL, pGAPZMFMHDEL, pPICZBMFMHDEL, pGAPZmMhDEL, pGAPZmMhDEL; pPIC9mMhDEL and pGAPZmMhDEL; a method of reducing the glycosylation on proteins produced from a methylotrophic yeast, comprising transforming said yeast with said vectors; said method wherein said yeast is *Pichia pastoris*;

2. Claims: (11-19)-complete, (22-24)-partially

A vector capable of expressing a glucosidase II or a functional part thereof in a methylotrophic yeast strain, comprising a nucleotide sequence coding for said alpha-1,2-mannosidase or said functional part; a vector selected from the group consisting of pGAPZAGLSII, PAOX2ZAGLSII, pYPTIZAGLSII, pGAPADEglsII, pPICADEglsII, PAOX2ADEglsII, pYPTIADEglsII, pGAPZAglsIIHDEL and pGAPADEglsIIHDEL; a method of reducing the glycosylation on proteins produced from a methylotrophic yeast, comprising transforming said yeast with said vectors; said method wherein said yeast is *Pichia pastoris*;

3. Claims: (20-21)-complete, (22-24)-partially

A vector for disrupting the *Och1* gene in a methylotrophic yeast strain, comprising a portion of the *och1* gene and a selectable marker gene, wherein said portion of the *Och1* gene and said selectable marker gene are linked in such a way to effect the disruption of the genomic *Och1* gene in said methylotrophic yeast strain; a vector having the designation pBLURA5'PpOCH1; a method of reducing the glycosylation on proteins produced from a methylotrophic yeast, comprising transforming said yeast with said vectors; said method wherein said yeast is *Pichia pastoris*;

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 01/01514

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0548012	A	23-06-1993	AT 157703 T	15-09-1997
			AU 667852 B2	18-04-1996
			AU 3007192 A	17-06-1993
			CA 2085308 A1	17-06-1993
			DE 69222013 D1	09-10-1997
			DE 69222013 T2	29-01-1998
			DK 548012 T3	14-04-1998
			EP 0548012 A1	23-06-1993
			ES 2107520 T3	01-12-1997
			FI 925678 A	17-06-1993
			GR 3025324 T3	27-02-1998
			JP 3305781 B2	24-07-2002
			JP 6197787 A	19-07-1994
			KR 270283 B1	01-11-2000
			MX 9207291 A1	01-07-1993
			NO 924847 A	17-06-1993
			NZ 245469 A	28-03-1995
			US 5501975 A	26-03-1996
			US 5618690 A	08-04-1997
			ZA 9209711 A	16-06-1993
EP 1211310	A	05-06-2002	AU 6592800 A	19-03-2001
			EP 1211310 A1	05-06-2002
			WO 0114522 A1	01-03-2001
WO 9105057	A	18-04-1991	WO 9105057 A1	18-04-1991
EP 0582244	A	09-02-1994	DE 4301932 A1	10-02-1994
			AT 207955 T	15-11-2001
			AU 657230 B2	02-03-1995
			AU 4435493 A	24-02-1994
			CA 2103522 A1	08-02-1994
			DE 59310233 D1	06-12-2001
			EP 0582244 A2	09-02-1994
			FI 933487 A	08-02-1994
			JP 2984229 B2	29-11-1999
			JP 10052277 A	24-02-1998
			JP 2675511 B2	12-11-1997
			JP 6296482 A	25-10-1994
			KR 140973 B1	15-06-1998
			NO 932811 A	08-02-1994
			NZ 248323 A	22-12-1994
			US 5705616 A	06-01-1998
			US 5798226 A	25-08-1998
			ZA 9305719 A	06-02-1995
WO 9621038	A	11-07-1996	US 5834251 A	10-11-1998
			AU 4348696 A	24-07-1996
			EP 0799318 A1	08-10-1997
			WO 9621038 A1	11-07-1996
EP 0314096	A	03-05-1989	AT 100142 T	15-01-1994
			AU 2435688 A	11-05-1989
			CA 1325781 A1	04-01-1994
			DE 3887082 D1	24-02-1994
			DE 3887082 T2	28-04-1994
			DK 599788 A	30-04-1989
			EP 0314096 A2	03-05-1989

INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/IB 01/01514

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0314096	A	FI 884991 A	30-04-1989
		JP 2000419 A	05-01-1990
		US 5135854 A	04-08-1992